For Research Use Only

UNC5C-Specific Polyclonal antibody

Catalog Number: 20240-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

NM 003728

Size: GeneID (NCBI):

150ul , Concentration: 500 µg/ml by 8633

Nanodrop and 327 µg/ml by Bradford Full Name: method using BSA as the standard;

nethod using BSA as the standard; unc-5 homolog C (C. elegans)

Source: Calculated MW:
Rabbit 103 kDa
Isotype: Observed MW:
IgG 50 kDa, 103 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:8000

WB 1:1000-1:8000 IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

20240-1-AP

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, mouse kidney tissue, rat brain

tissue

IHC: mouse brain tissue,

Background Information

UNC5C, also named as UNC5H3, belongs to the unc-5 family. It is a receptor for netrin required for axon guidance. It mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. It also involved in corticospinal tract axon guidances independently of DCC. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. The antibody is specific to UNC5C.

Notable Publications

AuthorPubmed IDJournalApplicationShaista Chaudhary36940724Alcohol Clin Exp ResWB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20240-1-AP (UNC 5C-Specific antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20240-1-AP (UNC5C-Specific antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20240-1-AP (UNC5C-Specific antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).