For Research Use Only

INSR/CD220 Polyclonal antibody

Catalog Number:20433-1-AP 31 Publications



Basic Information

Catalog Number:

BC 117172

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 500 ug/ml by

20433-1-AP

WB 1:500-1:2000

Nanodrop:

UNIPROT ID: P06213

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Rabbit

Full Name: INSR

Isotype: IgG

AG13922

Calculated MW:

Immunogen Catalog Number:

1382 aa, 156 kDa

Observed MW:

190 kDa, 135 kDa, 95 kDa

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IHC, IF, CoIP, Dot blot

Species Specificity:

human

Cited Species:

human, mouse, rat, pig, zebrafish, fish

Positive Controls:

WB: HEK-293 cells, MCF-7 cells, HT-1080 cells, HeLa

cells

IP: HeLa cells,

Background Information

The biological effects of INS are mediated by a membrane-spanning cell surface receptor that has been shown to consist of disulfide-linked alpha-subunits (135 kDa) and beta-subunits (95 kDa) that are cleaved products of a common precursor. The binding of INS to the receptor extracellular domain results in intracellular activation of a tyrosine-specific kinase activity and the generation of signals that determine the cellular response (PMID: 2369896). This antibody raised against 980-1382aa of human INS receptor can recognize full-length INS receptor precursor and INS receptor beta subunit.

Notable Publications

Author	Pubmed ID	Journal	Application
Takashi Hara	36130217	Biosci Biotechnol Biochem	WB
Yu Bai	34603025	Front Pharmacol	WB,CoIP
Si-Lan Han	33245450	Fish Physiol Biochem	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data







HEK-293 cells were subjected to SDS PAGE followed by western blot with 20433-1-AP (INSR antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.