For Research Use Only

DYKDDDDK tag Polyclonal antibody (Binds to FLAG® tag epitope)

proteintech®

Antibodies | ELISA kits | Proteins

www.ptglab.com

Catalog Number: 20543-1-AP

1122 Publications

Basic Information

Catalog Number:

20543-1-AP

Size:

150ul , Concentration: 600 ug/ml by

Nanodrop;

Source: Rabbit Isotype: IgG

Immunogen Catalog Number:

AG2329

GenBank Accession Number:

GeneID (NCBI):

Full Name: Flag Tag Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:20000-1:100000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP, RIP

Species Specificity: recombinant protein Cited Species: human, mouse, pig

Positive Controls:

WB: Transfected HEK-293T cells, IP: Transfected HEK-293 cells,

Background Information

Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. The DYKDDDDK(FLAG) peptide has been used extensively as a general tag in expression vectors. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. N-terminal DDDDK vectors provide an Ek cleavage site for removal of the fusion tag. The DDDDK peptide is likely to be located on the surface of a fusion protein because of its hydrophilic nature. As a result, the DDDDK peptide is more likely to be accessible to antibodies. A DDDDK-tag can be used in many different assays that require recognition by an antibody, such as western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions, cell ultrastructure, and protein localization and so on. This antibody is a rabbit polyclonal antibody raised against 3xFlag (3xDYKDDDDKT) sequence and recognizes the (1x) and (3x)DYKDDDDK peptide and detects DDDDK-tagged proteins. Anti-FLAG is a registered trademark of Sigma-Aldrich Biotechnology.

Notable Publications

Author	Pubmed ID	Journal	Application
Sirwan Sleman	36179070	Viral Immunol	WB,IP
Huanru Wang	31575039	Int J Mol Sci	
M Zatyka	25274773	Hum Mol Genet	IP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



HEK-293T cells and transfected HEK-293T lysates were subjected to SDS PAGE followed by western blot with 20543-1-AP (DYKDDDDK tag antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



IP result of anti-DYKDDDDK tag (IP:20543-1-AP, 4ug; Detection:20543-1-AP 1:10000) with Transfected HEK-293 cells lysate 400 ug.