

For Research Use Only

CHD1 Polyclonal antibody

Catalog Number: 20576-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: 20576-1-AP	GenBank Accession Number: NM_001270	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop and 327 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 1105	Recommended Dilutions: WB 1:300-1:1000 IP 0.5-4.0 ug for IP and 1:200-1:1000 for WB
Source: Rabbit	Full Name: chromodomain helicase DNA binding protein 1	
Isotype: IgG	Calculated MW: 197 kDa	
	Observed MW: 200 kDa	

Applications

Tested Applications: IP, WB, ELISA	Positive Controls: WB : HeLa cells, A431 cells, human brain tissue, Jurkat cells IP : HeLa cells,
Cited Applications: IF, WB	
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Background Information

CHD1 belongs to the SNF2/RAD54 helicase family. CHD1 is an ATP-dependent chromatin-remodeling factor which functions as substrate recognition component of the transcription regulatory histone acetylation (HAT) complex SAGA. CHD1 regulates polymerase II transcription. It is required for efficient transcription by RNA polymerase I, and more specifically the polymerase I transcription termination step. CHD1 regulates negatively DNA replication. It is required to maintain a specific chromatin configuration across the genome. It is associated with histone deacetylase (HDAC) activity. CHD1 is required for the bridging of SNF2, the FACT complex, the PAF complex as well as the U2 snRNP complex to H3K4me3. It functions to modulate the efficiency of pre-mRNA splicing in part through physical bridging of spliceosomal components to H3K4me3. It is required for maintaining open chromatin and pluripotency in embryonic stem cells. In PTEN-deficient prostate and breast cancers, CHD1 depletion profoundly and specifically suppressed cell proliferation, cell survival and tumorigenic potential. The antibody is specific to CHD1.

Notable Publications

Author	Pubmed ID	Journal	Application
Jun Zhu	31667976	Cell Prolif	WB
Peter Rüthemann	29018037	EMBO J	WB
Alexandra Wille	26635563	Front Behav Neurosci	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 20576-1-AP (CHD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.

IP Result of anti-CHD1 (IP:20576-1-AP, 4ug; Detection:20576-1-AP 1:300) with HeLa cells lysate 1600ug.