For Research Use Only

CHD1 Polyclonal antibody

Catalog Number:20576-1-AP

Featured Product

6 Publications





Basic Information	Catalog Number: 20576-1-AP	GenBank Accession Number: NM_001270	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 700 ug/ml by	1105	WB 1:300-1:1000	
	Nanodrop and 327 ug/ml by Bradford method using BSA as the standard; Source:	UNIPROT ID: 014646	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
		Full Name:		
	Rabbit	chromodomain helicase DNA binding protein 1		
	Isotype: IgG			
		Calculated MW: 197 kDa		
		Observed MW: 200 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IP, ELISA Cited Applications:		WB : HeLa cells, A431 cells, human brain tissue, Jurka	
	WB, IF	cells		
	Species Specificity:			
	human, mouse, rat			
	Cited Species:			
	human, mouse			
Background Information	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor	nponent of the transcription regu	latory histone acetylation (HAT) complex	
Background Information	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor SAGA. CHD1 regulates polymerase I more specifically the polymerase I tr required to maintain a specific chrom (HDAC) activity. CHD1 is required for snRNP complex to H3K4me3. It functi bridging of spliceosomal components	nponent of the transcription regu transcription. It is required for ef anscription termination step. CHI latin configuration across the ger the bridging of SNF2, the FACT c ons to modulate the efficiency of s to H3K4me3. It is required for m cient prostate and breast cancers	ficient transcription by RNA polymerase I, an D1 regulates negatively DNA replication. It is nome. It is associated with histone deacetylas complex, the PAF complex as well as the U2 f pre-mRNA splicing in part through physical naintaining open chromatin and pluripotency s, CHD1 depletion profoundly and specifically	
	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor SAGA. CHD1 regulates polymerase II more specifically the polymerase I tr required to maintain a specific chrom (HDAC) activity. CHD1 is required for snRNP complex to H3K4me3. It functi bridging of spliceosomal components in embryonic stem cells. In PTEN-defi suppressed cell proliferation, cell sur	nponent of the transcription regu transcription. It is required for ef anscription termination step. CHI latin configuration across the ger the bridging of SNF2, the FACT c ons to modulate the efficiency of s to H3K4me3. It is required for m cient prostate and breast cancers	latory histone acetylation (HAT) complex ficient transcription by RNA polymerase I, an D1 regulates negatively DNA replication. It is nome. It is associated with histone deacetylas complex, the PAF complex as well as the U2 f pre-mRNA splicing in part through physical naintaining open chromatin and pluripotency s, CHD1 depletion profoundly and specifically	
	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor SAGA. CHD1 regulates polymerase II more specifically the polymerase I to required to maintain a specific chrom (HDAC) activity. CHD1 is required for snRNP complex to H3K4me3. It functi bridging of spliceosomal components in embryonic stem cells. In PTEN-defi suppressed cell proliferation, cell sur	nponent of the transcription regu transcription. It is required for ef anscription termination step. CHI latin configuration across the ger the bridging of SNF2, the FACT c ons to modulate the efficiency of s to H3K4me3. It is required for m cient prostate and breast cancers vival and tumorigenic potential.	latory histone acetylation (HAT) complex ficient transcription by RNA polymerase I, an D1 regulates negatively DNA replication. It is nome. It is associated with histone deacetylas complex, the PAF complex as well as the U2 f pre-mRNA splicing in part through physical naintaining open chromatin and pluripotency s, CHD1 depletion profoundly and specifically The antibody is specific to CHD1.	
Background Information	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor SAGA. CHD1 regulates polymerase I I more specifically the polymerase I tr required to maintain a specific chrom (HDAC) activity. CHD1 is required for snRNP complex to H3K4me3. It functi bridging of spliceosomal components in embryonic stem cells. In PTEN-defi suppressed cell proliferation, cell sur Author Pul Jun Zhu 31	nponent of the transcription regu transcription. It is required for ef anscription termination step. CH atin configuration across the ger the bridging of SNF2, the FACT c ons to modulate the efficiency of is to H3K4me3. It is required for m cient prostate and breast cancers vival and tumorigenic potential.	Idatory histone acetylation (HAT) complex ficient transcription by RNA polymerase I, an D1 regulates negatively DNA replication. It is nome. It is associated with histone deacetylas complex, the PAF complex as well as the U2 f pre-mRNA splicing in part through physical naintaining open chromatin and pluripotency s, CHD1 depletion profoundly and specifically The antibody is specific to CHD1.	
	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor SAGA. CHD1 regulates polymerase II more specifically the polymerase I tr required to maintain a specific chrom (HDAC) activity. CHD1 is required for snRNP complex to H3K4me3. It functi bridging of spliceosomal components in embryonic stem cells. In PTEN-defi suppressed cell proliferation, cell sur Author Pul Jun Zhu 311 Peter Rüthemann 290	nponent of the transcription regularized for eff anscription. It is required for eff anscription termination step. CHI latin configuration across the ger the bridging of SNF2, the FACT of ons to modulate the efficiency of to H3K4me3. It is required for m cient prostate and breast cancers vival and tumorigenic potential.	Idatory histone acetylation (HAT) complex ficient transcription by RNA polymerase I, an D1 regulates negatively DNA replication. It is nome. It is associated with histone deacetyla complex, the PAF complex as well as the U2 f pre-mRNA splicing in part through physical haintaining open chromatin and pluripotency s, CHD1 depletion profoundly and specifically The antibody is specific to CHD1. Application WB WB	
	CHD1 belongs to the SNF2/RAD54 he functions as substrate recognition cor SAGA. CHD1 regulates polymerase II more specifically the polymerase I tr required to maintain a specific chrom (HDAC) activity. CHD1 is required for snRNP complex to H3K4me3. It functi bridging of spliceosomal components in embryonic stem cells. In PTEN-defi suppressed cell proliferation, cell sur Author Pul Jun Zhu 311 Peter Rüthemann 290	nponent of the transcription regularies transcription. It is required for effanscription termination step. CHI and the configuration across the gere the bridging of SNF2, the FACT of so to H3K4me3. It is required for micient prostate and breast cancers vival and tumorigenic potential.	Idatory histone acetylation (HAT) complex ficient transcription by RNA polymerase I, an D1 regulates negatively DNA replication. It is nome. It is associated with histone deacetylat complex, the PAF complex as well as the U2 f pre-mRNA splicing in part through physical vaintaining open chromatin and pluripotency s, CHD1 depletion profoundly and specifically The antibody is specific to CHD1. Application WB WB	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





HeLa cells were subjected to SDS PAGE followed by western blot with 20576-1-AP (CHD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. IP result of anti-CHD1 (IP:20576-1-AP, 4ug; Detection:20576-1-AP 1:300) with HeLa cells lysate 1600ug.