For Research Use Only

## AIM2 Polyclonal antibody

Catalog Number:20590-1-AP

Featured Product






Basic Information	Catalog Number: 20590-1-AP	GenBank Accession Number: BC010940 GeneID (NCBI): 9447 UNIPROT ID: 014862 Full Name:		Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000 IHC 1:20-1:200 IF/ICC 1:200-1:800		
	Size:					
	150ul , Concentration: 750 ug/ml by					
	Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG14520					
				absent in melanoma 2	2	
		Calculated MW:				
		343 aa, 39 kDa				
		Observed MW: 39-42 kDa				
		Applications	Tested Applications:		Positive Controls:	
WB, IHC, ELISA			WB : A431 cells, HeLa cells, Jurkat cells, SH-SY5Y cells			
Cited Applications: WB, IHC	mouse testis tissue, rat testis tissue, MCF-7 cells, A54 cells, COLO 320 cells, HepG2 cells, Raji cells					
Species Specificity:			IHC : human testis tissue,			
human, mouse, rat			IF/ICC : MCF-7 cells,			
Cited Species: human, mouse, rat, zebrafish						
Note-IHC: suggested antigen ( TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen					
	Absent in melanoma 2 (AIM2) is a se and host immune responses to DNA	/iruses and intracellular	bacteria [PMI	D:22695634]. Upon b		
Background Information	thought to undergo oligomerization and processing of interleukin-1 beta by repressing NF-kappa-B transcripti	and interleukin-18. Oth	erwise, it funct	tions as a tumor supp	caspase-1 precusrsor	
	thought to undergo ol igomerization and processing of interleukin-1 beta by repressing NF-kappa-B transcripti	and interleukin-18. Oth	erwise, it funct 32157,2056683	tions as a tumor supp	caspase-1 precusrsor	
	thought to undergo oligomerization and processing of interleukin-1 beta by repressing NF-kappa-B transcripti Author Pu	and interleukin-18. Oth onal activity [PMID:164] bmed ID Journ	erwise, it funct 32157,2056683	tions as a tumor supp	caspase-1 precusrsor pressor which may ac	
Background Information Notable Publications	thought to undergo oligomerization and processing of interleukin-1 beta by repressing NF-kappa-B transcripti Author Pu Liangmin Fu 36	and interleukin-18. Oth onal activity [PMID:164] bmed ID Journ 248876 Front	erwise, it funct 32157,2056683 al	tions as a tumor supp	caspase-1 precusrsor pressor which may ac Application	
	thought to undergo oligomerization and processing of interleukin-1 beta by repressing NF-kappa-B transcripti Author Pu Liangmin Fu 36 JiangFan Yu 36	and interleukin-18. Oth onal activity [PMID:164] bmed ID Journ 248876 Front 248980 Front	erwise, it funct 32157,2056683 al Immunol	tions as a tumor sup 31].	caspase-1 precusrso pressor which may ac Application WB,IHC	
	thought to undergo oligomerization and processing of interleukin-1 beta by repressing NF-kappa-B transcripti Author Pu Liangmin Fu 36 JiangFan Yu 36	and interleukin-18. Oth onal activity [PMID:164] bmed ID Journ 248876 Front 248980 Front 037355 Proc N ter shipment.	erwise, it funct 32157,205668 al Immunol Oncol	tions as a tumor sup 31].	caspase-1 precusrso pressor which may ac Application WB,IHC WB	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20590-1-AP (AIM2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 20590-1-AP (AIM2 antibody) at dilution of 1:100 (under 10x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 20590-1-AP (AIM2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using AIM2 antibody (20590-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).