For Research Use Only

MARCKS Polyclonal antibody

Catalog Number: 20661-1-AP

3 Publications



Basic Information

Catalog Number: GenBank Accession Number: 20661-1-AP

GeneID (NCBI):

150ul, Concentration: 550 µg/ml by 4082

Nanodrop;

Source: myristoylated alanine-rich protein

Rabbit kinase C substrate Isotype: Calculated MW: 32 kDa IgG

> Observed MW: 80 kDa

NM 002356

Positive Controls:

WB: HEK-293 cells, mouse brain tissue, rat brain tissue

Purification Method:

WB 1:500-1:2000

IHC 1:50-1:500

IF 1:200-1:800

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:200-1:1000

Recommended Dilutions:

IP: A431 cells,

IHC: mouse brain tissue, IF: mouse brain tissue,

Applications

Tested Applications: IF, IHC, IP, WB, ELISA

Cited Applications:

Species Specificity: human, mouse, rat **Cited Species:**

human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

MARCKS, also named as MACS, PRKCSL and 80K-L protein, belongs to the MARCKS family. MARCKS is the most prominent cellular substrate for protein kinase C. It binds calmodulin, actin, and synapsin. MARCKS is a filamentous (F) actin cross-linking protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Nicholas J Eustace	30942445	Int J Oncol	IP
Falguni Das	35835217	J Biol Chem	WB
Tong Lei	32800794	J Proteomics	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20661-1-AP (MARCKS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20661-1-AP (MARCKS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using MARCKS antibody (20661-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP Result of anti-MARCKS (IP:20661-1-AP, 4ug; Detection:20661-1-AP 1:300) with A431 cells lysate 1200ug.