

For Research Use Only

PI3 Kinase p110 Gamma Polyclonal antibody



Catalog Number: 20662-1-AP

4 Publications

Basic Information

Catalog Number:

20662-1-AP

Size:

150ul, Concentration: 700 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_002649

GeneID (NCBI):

5294

Full Name:

phosphoinositide-3-kinase, catalytic, gamma polypeptide

Calculated MW:

126 kDa

Observed MW:

100-130 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 µg for IP and 1:200-1:1000

for WB

IHC 1:200-1:800

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

Rat, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse skeletal muscle tissue, rat skeletal muscle tissue

IP: mouse liver tissue,

IHC: mouse skeletal muscle tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Shuo Chen	36056690	J Cell Mol Med	WB
Xiaoyan Wan	32603600	Am J Physiol Endocrinol Metab	WB
Yi Hou	35834807	J Med Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20662-1-AP (PI3 Kinase p110 Gamma antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP Result of anti-PI3K p110(gamma) (IP:20662-1-AP, 5ug; Detection:20662-1-AP 1:300) with mouse liver tissue lysate 4000ug.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 20662-1-AP (PI3 Kinase p110 Gamma antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with .



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 20662-1-AP (PI3 Kinase p110 Gamma antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with .