For Research Use Only

IRS2 Polyclonal antibody Catalog Number: 20702-1-AP 11 Publications




Basic Information	Catalog Number: 20702-1-AP	GenBank Accession Number: NM_003749	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 500 ug/ml by	8660	WB 1:500-1:1000	
	Nanodrop;	UNIPROT ID:	IHC 1:50-1:500	
	Source: Rabbit	Q9Y4H2		
	Isotype:	Full Name: IRS2		
	lgG	Calculated MW: 137 kDa		
		Observed MW: 150-185 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, ELISA	WB : mo	WB : mouse brain tissue,	
	Cited Applications: WB, IF	IHC : mouse skeletal muscle tissue,		
	Species Specificity: human, mouse			
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen		
	The family of INS receptor substrates (IRS) consists of four proteins (IRS-1-IRS-4), which were initially characterized as typical cytosolic adaptor proteins involved in INS receptor (IR) and INS-like growth factor I receptor (IGF-IR) signaling. Impaired glucose tolerance and INS resistance, and thus type 2 diabetes were demonstrated in Irs2 knockout mice. In vivo data also indicated that he regulation of beta-cell mass involves IRS2, while IRS1 is only required for proper INS production in beta-cells. IRS2, like IRS1, has a predicted molecular weight of 137 kDa, however, as a result of its extensive serine phosphorylation it separates on a SDS-PAGE as a band of approximately 150-185 kDa.			
Background Information	as typical cytosolic adaptor proteins signaling. Impaired glucose toleranc knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa,	
	as typical cytosolic adaptor proteins signaling. Impaired glucose toleranc knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive 150-185 kDa.	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa, tes on a SDS-PAGE as a band of approximate	
	as typical cytosolic adaptor proteins signaling. Impaired glucose toleranc knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive 150-185 kDa.	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa med ID Journal	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa, tes on a SDS-PAGE as a band of approximate Application	
	as typical cytosolic adaptor proteins signaling. Impaired glucose toleranc knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive 150-185 kDa. Author Put Li Lin 363	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa omed ID Journal 525267 Genet Res (Can	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa, tes on a SDS-PAGE as a band of approximate Application mb) WB	
Background Information	as typical cytosolic adaptor proteins signaling. Impaired glucose toleranc knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive 150-185 kDa. Author Put Li Lin 363	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa med ID Journal	INS-like growth factor I receptor (IGF-I pe 2 diabetes were demonstrated in Irs cell mass involves IRS2, while IRS1 is of predicted molecular weight of 137 kDa tes on a SDS-PAGE as a band of approxi Application nb) WB	
Notable Publications	as typical cytosolic adaptor proteins signaling. Impaired glucose tolerance knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive so 150-185 kDa. Author Put Li Lin 362 Hang Peng 358 Yuhua Xu 350	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa omed ID Journal 525267 Genet Res (Can	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa, tes on a SDS-PAGE as a band of approximation mb) WB WB,IF	
Notable Publications	as typical cytosolic adaptor proteins signaling. Impaired glucose tolerance knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive so 150-185 kDa. Author Put Li Lin 363 Hang Peng 358 Yuhua Xu 350 Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa omed ID Journal 325267 Genet Res (Can 344799 Int J Biol Sci 388561 Obesity (Silver er shipment.	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa, tes on a SDS-PAGE as a band of approximate hb) WB WB,IF	
	as typical cytosolic adaptor proteins signaling. Impaired glucose toleranc knockout mice. In vivo data also indi required for proper INS production in however, as a result of its extensive s 150-185 kDa. Author Put Li Lin 362 Hang Peng 356 Yuhua Xu 350 Storage: Stora et -20°C. Stable for one year aft Storage Buffer:	e and INS resistance, and thus ty cated that he regulation of beta- beta-cells. IRS2, like IRS1, has a serine phosphorylation it separa omed ID Journal 325267 Genet Res (Can 344799 Int J Biol Sci 388561 Obesity (Silver er shipment.	INS-like growth factor I receptor (IGF-IR) pe 2 diabetes were demonstrated in Irs2 cell mass involves IRS2, while IRS1 is only predicted molecular weight of 137 kDa, tes on a SDS-PAGE as a band of approximate hb) WB WB,IF	

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse brain tissue were subjected to SDS PAGE followed by western blot with 20702-1-AP (IRS2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 20702-1-AP (IRS2 antibody) at dilution of 1:200 (under 10x lens).