For Research Use Only

GAD65 Polyclonal antibody

Catalog Number:20746-1-AP 20 Publications



Basic Information

Catalog Number:

20746-1-AP

Size:

150ul , Concentration: 750 ug/ml by 2572
Nanodrop and 547 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; 005320

Source:

Rabbit Isotype:

IgG

GenBank Accession Number:

NM_000818 GeneID (NCBI):

UNIPROT ID: Q05329 Full Name:

glutamate decarboxylase 2 (pancreatic islets and brain, 65kDa)

Calculated MW: 65 kDa Observed MW: 65 kDa Purification Method:

Antigen affinity purification
Recommended Dilutions:
WB 1:3000-1:20000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications: WB, IP, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat Positive Controls:

WB: mouse brain tissue, rat brain tissue

IP: mouse brain tissue,

Background Information

GAD2, also named as GAD65, belongs to the group II decarboxylase family. GAD2 catalyzes the production of GABA. It is responsible for the synthesis of the essential neurotransmitter gamma-aminobutyric acid (GABA) from L-glutamic acid. GAD2 is expressed in nervous and endocrine systems and are thought to be involved in synaptic transmission and INS secretion. Autoantibodies against GAD2 may serve as markers for type I diabetes. Many individuals suffering from an adult onset disorder known as Stiff Person Syndrome (SPS) also express autoantibodies to GAD2. The antibody is specific to GAD2.

Notable Publications

Author	Pubmed ID	Journal	Application
Yueqin Liu	33344711	Neurobiol Stress	WB
Hongru Yang	35618610	Adv Sci (Weinh)	WB
Ken Chen	31015358	Clin Sci (Lond)	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20746-1-AP (GAD65 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-GAD65 (IP:20746-1-AP, 4ug; Detection:20746-1-AP 1:1000) with mouse brain tissue lysate 3440ug.