

For Research Use Only

ZNF397 Polyclonal antibody

Catalog Number: 20799-1-AP



Basic Information

Catalog Number: 20799-1-AP	GenBank Accession Number: BC006172	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 1200 ug/ml by Nanodrop and 593 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 84307	Recommended Dilutions: WB 1:200-1:1000 IHC 1:250-1:1000
Source: Rabbit	UNIPROT ID: Q8NF99	
Isotype: IgG	Full Name: zinc finger protein 397	
Immunogen Catalog Number: AG14712	Calculated MW: 534 aa, 61 kDa	
	Observed MW: 60-63 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Species Specificity: human, mouse, Rat	WB : PC-3 cells, mouse testis tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse liver tissue,

Background Information

ZNF397 (Zinc Finger Protein 397) is conserved in placental mammals. ZNF397 as a bona fide coactivator of the androgen receptor (AR), essential for the transcriptional program governing AR-driven luminal lineage.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 20799-1-AP (ZNF397 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 20799-1-AP (ZNF397 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



PC-3 cells were subjected to SDS PAGE followed by western blot with 20799-1-AP (ZNF397 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.