For Research Use Only

## TBR1 Polyclonal antibody Catalog Number: 20932-1-AP 35 Publications






| Basic Information                              | Catalog Number:<br>20932-1-AP   | GenBank Accession Number:<br>BC 104844  |   | Purification Method:<br>Antigen affinity purification  |           |  |
|--|---|---|---|--|-----------|--|
|  | Size:<br>150ul, Concentration: 750 ug/ml by<br>Nanodrop;<br>Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG14935  | GeneID (NCBI):<br>10716<br>UNIPROT ID:<br>Q16650<br>Full Name:<br>T-box, brain, 1<br>Calculated MW:<br>682 aa, 74 kDa<br>Observed MW:<br>74 kDa |   | Recommended Dilutions:<br>WB 1:500-1:1000<br>IP 0.5-4.0 ug for 1.0-3.0 mg of<br>protein lysate<br>IHC 1:20-1:200<br>IF-P 1:200-1:800 | total     |  |
| Applications                                   | Tested Applications:  |   | Positive Controls:  |  |           |  |
|  | WB, IHC, IF-P, IP, ELISA<br>Cited Applications:   |   | WB : mouse brain tissue,  |  |           |  |
|  | WB, IHC, IF   |   |   |  |           |  |
|  | Species Specificity:<br>human, mouse, rat   |   | IHC : human l<br>tissue   | IC : human brain tissue, mouse brain tissue, rat bra<br>ssue   |           |  |
|  | Cited Species:<br>human, mouse, rat   |   | IF-P : mouse brain tissue,  |  |           |  |
|  | Note-IHC: suggested antigen (<br>TE buffer pH 9.0; (*) Alternati<br>retrieval may be performed w<br>buffer pH 6.0   | vely, antigen   |   |  |           |  |
|  |   |   |   |  |           |  |
| Background Information                         | TBR1, also named as T-box brain pro<br>domain and localizes in the nucleus.<br>developmental processes. TBR1 is re  | TBR1 is expressed in t  | he brain and as   |  | -         |  |
|  | domain and localizes in the nucleus.<br>developmental processes. TBR1 is re   | TBR1 is expressed in t<br>quired for normal brai  | he brain and as<br>n development.                                     | a transcriptional regulator is ir  | volved in |  |
|  | domain and localizes in the nucleus.<br>developmental processes. TBR1 is re<br>Author Pr  | TBR1 is expressed in t<br>quired for normal brai<br>ubmed ID Jou  | he brain and as<br>n development.<br>rnal                             | a transcriptional regulator is ir<br>Applica   | volved in |  |
| Background Information<br>Notable Publications | domain and localizes in the nucleus.<br>developmental processes. TBR1 is re<br>Author Pu<br>Inseyah Bagasrawala 27  | TBR1 is expressed in t<br>quired for normal brai<br>ubmed ID Jou<br>1746712 Fro   | he brain and as<br>n development.                                     | a transcriptional regulator is in<br>Applica<br>WB,IF  | volved in |  |
|  | domain and localizes in the nucleus.<br>developmental processes. TBR1 is reAuthorPtInseyah Bagasrawala27Inseyah Bagasrawala27   | TBR1 is expressed in t<br>quired for normal brai<br><b>Ibmed ID Jou</b><br>1746712 From<br>1664962 Cer  | he brain and as<br>n development.<br>rnal<br>nt Neurosci              | a transcriptional regulator is ir<br>Applica   | volved in |  |
|  | domain and localizes in the nucleus.<br>developmental processes. TBR1 is reAuthorPtInseyah Bagasrawala27Inseyah Bagasrawala27   | TBR1 is expressed in t<br>quired for normal brai<br>bmed ID Jou<br>7746712 From<br>7664962 Cer<br>0307948 PLo<br>er shipment.                   | he brain and as<br>n development.<br>rnal<br>nt Neurosci<br>eb Cortex | a transcriptional regulator is in<br>Applica<br>WB,IF<br>WB,IHC  | volved in |  |
| Notable Publications                           | domain and localizes in the nucleus.   developmental processes. TBR1 is re   Author Pet   Inseyah Bagasrawala 27   Inseyah Bagasrawala 27   Giedre Miskinyte 36   Storage: Storage Buffer:   PBS with 0.02% sodium azide and 50 | TBR1 is expressed in t<br>quired for normal brai<br>bmed ID Jou<br>7746712 From<br>7664962 Cer<br>0307948 PLo<br>er shipment.                   | he brain and as<br>n development.<br>rnal<br>nt Neurosci<br>eb Cortex | a transcriptional regulator is in<br>Applica<br>WB,IF<br>WB,IHC  | volved in |  |
| Notable Publications                           | domain and localizes in the nucleus.   developmental processes. TBR1 is re   Author Pet   Inseyah Bagasrawala 27   Inseyah Bagasrawala 27   Giedre Miskinyte 36   Storage: Storage Buffer:   PBS with 0.02% sodium azide and 50 | TBR1 is expressed in t<br>quired for normal brai<br>bmed ID Jou<br>7746712 From<br>7664962 Cer<br>0307948 PLo<br>er shipment.                   | he brain and as<br>n development.<br>rnal<br>nt Neurosci<br>eb Cortex | a transcriptional regulator is in<br>Applica<br>WB,IF<br>WB,IHC  | volved in |  |


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using TBR1 antibody (20932-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




mouse brain tissue were subjected to SDS PAGE followed by western blot with 20932-1-AP (TBR1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human brain using 20932-1-AP (TBR1 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human brain using 20932-1-AP (TBR1 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-TBR1 (IP:20932-1-AP, 4ug; Detection:20932-1-AP 1:600) with mouse brain tissue lysate 4000ug.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using TBR1 antibody (20932-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).