

For Research Use Only

HIF-1 alpha Polyclonal antibody

Catalog Number: 20960-1-AP

Featured Product

511 Publications



Basic Information

Catalog Number:

20960-1-AP

Size:

150ul, Concentration: 600 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG15198

GenBank Accession Number:

BC012527

GeneID (NCBI):

3091

Full Name:

hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)

Calculated MW:

826 aa, 93 kDa

Observed MW:

120 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 µg for 1.0-3.0 mg of total

protein lysate

IHC 1:50-1:500

IF 1:200-1:800

Applications

Tested Applications:

FC, IF, IHC, IP, WB, ELISA

Cited Applications:

Cell treatment, ChIP, CoIP, IF, IHC, IP, WB

Species Specificity:

human

Cited Species:

human, goat, chicken, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Cobalt Chloride treated HeLa cells, HeLa cells

IP: HeLa cells,

IHC: human thyroid cancer tissue, human heart tissue, human kidney tissue

IF: Cobalt Chloride treated HeLa cells, Cobalt Chloride treated HepG2 cells

Background Information

HIF1α, the major regulator of the cellular responses to hypoxia, consists of an oxygen-sensitive subunit, HIF1α (HIF1A), and an oxygen-insensitive subunit, HIF1β (arylhydrocarbon receptor nuclear transporter [ARNT]). Under normal oxygen conditions, HIF1α is continuously produced and destroyed, in a process involving hydroxylation, interaction with von Hippel-Lindau (VHL) protein, polyubiquitylation and subsequent proteasomal degradation. Under hypoxic conditions, hydroxylation is impaired and HIF1α is stabilized. HIF1α localizes in cytoplasm in normoxia, but it can translocate into nuclear in response to hypoxia. The calculated molecular weight of HIF1α is 93 kDa, but the modified protein HIF1α is about 110-120kDa (PMID: 11698256, PMID: 7539918).

Notable Publications

Author	Pubmed ID	Journal	Application
Menghui Xu	36290677	Antioxidants (Basel)	WB
Di Cui	34586717	Adv Healthc Mater	WB, IHC
Wenmin Yu	34676302	Open Med (Wars)	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using 20960-1-AP (HIF1a antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of HIF-1 alpha antibody (20960-1-AP; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HIF-1 alpha transfected HeLa cells. Sample 1: non-treated sh-Control transfected HeLa cells, Sample 2: Cobalt Chloride treated sh-Control transfected HeLa cells, Sample 3: Cobalt Chloride treated sh-HIF-1 alpha transfected HeLa cells.




Untreated and Cobalt Chloride treated HeLa cells were subjected to SDS PAGE followed by western blot with 20960-1-AP (HIF1a antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Cobalt Chloride treated HeLa cells using HIF-1 alpha antibody (20960-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



IP Result of anti-HIF1a (IP:20960-1-AP, 4ug; Detection:20960-1-AP 1:300) with HeLa cells lysate 4000ug.



1X10⁶ HeLa cells were stained with 0.2ug HIF1a antibody (20960-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.