For Research Use Only

MEKK3 Polyclonal antibody

Catalog Number:21072-1-AP

1 Publications



Purification Method:

WB 1:200-1:1000

IHC 1:50-1:500

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:200-1:1000

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

21072-1-AP BC090859 GeneID (NCBI):

150ul, Concentration: 800 µg/ml by 4215

Nanodrop;

Source: mitogen-activated protein kinase

Observed MW:

kinase kinase 3 Rabbit

Isotype: Calculated MW: 657 aa, 74 kDa IgG

Immunogen Catalog Number: 71 kDa, 74 kDa AG15223

Applications

Tested Applications:

IHC, IP, WB, ELISA Cited Applications:

IF, IP, WB

Species Specificity:

human

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse liver tissue, IP: mouse liver tissue,

IHC: human liver cancer tissue.

Background Information

The mitogen-activated protein/ERK kinase kinase 3 (MEKK3) is a member of a family of MEKK/Ste11 serine/threonine protein kinases. One function of MEKK3 is to serve as an upstream regulatory kinase in the $mitogen-activated\ protein\ kinase\ cascade,\ however,\ it\ also\ plays\ a\ critical\ role\ in\ other\ signaling\ pathways.\ MEKK3$ regulates several important functions including cell survival and proliferation. It is ubiquitously expressed in multiple tissues and has been found in complexes with scaffold proteins and other kinases. (PMID: 18206350). Cerebral cavernous malformations (CCMs) complex regulation of MEKK3 is essential during vertebrate heart development. MEKK3 signalling and KLF2/4 may be targeted to develop new CCM therapeutics.

Notable Publications

Application Author **Pubmed ID** Journal Jingiu Lu Oncogene WB,IF,IP 29662197

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




mouse liver tissue were subjected to SDS PAGE followed by western blot with 21072-1-AP (MEKK3 antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.



IP Result of anti-MEKK3 (IP:21072-1-AP, 4ug; Detection:21072-1-AP 1:300) with mouse liver tissue lysate 7000ug.

Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 21072-1-AP (MEKK3 antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 21072-1-AP (MEKK3 antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).