## For Research Use Only

## TBC1D19 Polyclonal antibody

Catalog Number: 21085-1-AP



**Purification Method:** 

WB 1:500-1:2000

IHC 1:50-1:500

IF 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number: 21085-1-AP BC031642

Size: GeneID (NCBI): 150ul, Concentration: 400 µg/ml by 55296

Nanodrop and 200 µg/ml by Bradford Full Name: method using BSA as the standard; TBC1 domain family, member 19

Calculated MW: Rabbit 526 aa, 60 kDa Isotype: Observed MW: IgG 55 kDa

Immunogen Catalog Number:

AG15323

**Positive Controls:** 

IF, IHC, WB, ELISA WB: mouse brain tissue, rat brain tissue Species Specificity: IHC: human pancreas cancer tissue, human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Tested Applications:** 

IF: HepG2 cells,

**Background Information** 

The function of TBC1D19 remains largely unknown. TBC1D19 may act as a GTPase-activating protein for Rab family protein(s).

Storage

**Applications** 

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



mouse brain tissue were subjected to SDS PAGE followed by western blot with 21085-1-AP (TBC 1D19 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immun of luorescent analysis of HepG2 cells using \$21085-1-AP (TBC 1D19 antibody)\$ at dilution of 1:50 and Alexa Fluor 488-conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 21085-1-AP (TBC1D19 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).