

For Research Use Only

C17orf28 Polyclonal antibody

Catalog Number: 21174-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number:
21174-1-AP

Size:
150ul, Concentration: 550 µg/ml by Nanodrop and 320 µg/ml by Bradford method using BSA as the standard;

Source:
Rabbit

Isotype:
IgG

Immunogen Catalog Number:
AG15427

GenBank Accession Number:
BC035372

GeneID (NCBI):
283987

Full Name:
chromosome 17 open reading frame 28

Calculated MW:
788 aa, 89 kDa

Observed MW:
60 kDa, 89 kDa

Purification Method:
Antigen affinity purification

Recommended Dilutions:
WB 1:500-1:2000
IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
IF 1:10-1:100

Applications

Tested Applications:
IF, IP, WB, ELISA

Cited Applications:
IF, WB

Species Specificity:
human, mouse, rat

Cited Species:
mouse, rat

Positive Controls:

WB: fetal human brain tissue, human brain tissue

IP: mouse brain tissue,

IF: HepG2 cells, Hela cells

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Wen Du	27751232	Elife	WB, IF
Min Li	30596141	Biophys Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



fetal human brain tissue were subjected to SDS PAGE followed by western blot with 21174-1-AP (C17orf28 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-C17orf28 (IP:21174-1-AP, 3ug; Detection:21174-1-AP 1:500) with mouse brain tissue lysate 10000ug.



Immunofluorescent analysis of HepG2 cells, using C17orf28 antibody 21174-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).