

For Research Use Only

MOGAT2 Polyclonal antibody

Catalog Number: 21286-1-AP



Basic Information

Catalog Number: 21286-1-AP	GenBank Accession Number: BC103878	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 133 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 80168	Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200
Source: Rabbit	Full Name: monoacylglycerol O-acyltransferase 2	
Isotype: IgG	Calculated MW: 334 aa, 38 kDa	
Immunogen Catalog Number: AG15806	Observed MW: 28-30 kDa	

Applications

Tested Applications:
IHC, WB, ELISA

Species Specificity:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HL-60 cells,

IHC : human kidney tissue, human stomach tissue

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HL-60 cells were subjected to SDS PAGE followed by western blot with 21286-1-AP (MOGAT2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney using 21286-1-AP (MOGAT2 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human kidney using 21286-1-AP (MOGAT2 antibody) at dilution of 1:50 (under 40x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 21286-1-AP (MOGAT2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.