## For Research Use Only

## Cathepsin D Polyclonal antibody, PBS Only



Catalog Number: 21327-1-PBS

**Featured Product** 

**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

Antigen affinity purification

21327-1-PBS

Nanodrop:

Isotype:

BC016320

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

**UNIPROT ID:** P07339

Rabbit

Full Name: cathepsin D

IgG Immunogen Catalog Number:

Calculated MW: 412 aa, 45 kDa

AG15254

Observed MW:

32 kDa, 48 kDa, 52 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse

**Background Information** 


CTSD (Cathepsin D) also named CPSD, belongs to the peptidase A1 family. It is ubiquitously expressed and is involved in proteolytic degradation, cell invasion, and apoptosis. Human CTSD is synthesized as a 52-kDa precursor that is converted into an active 48-kDa single-chain intermediate in the endosomes, and then into a fully active mature form, composed of a 34-kDa heavy chain and a 14-kDa light chain, in the lysosomes. It is a lysosomal acid protease found in neutrophils and monocytes and involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease (PMID: 27114232, PMID: 30717773, PMID: 30051532).

Storage

Store at -80°C. Storage Buffer:

PBS only, pH7.3


## Selected Validation Data



WB result of Cathepsin D antibody (21327-1-AP; 1:6000; room temperature for 1.5 hours) with wild-type and sg-Cathepsin D transfected HCT-116 cells. This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 21327-1-AP (Cathepsin D antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human hepatocirrhosis tissue slide using 21327-1-AP (Cathepsin D antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human hepatocirrhosis tissue slide using 21327-1-AP (Cathepsin D antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 21327-1-AP (Cathepsin D antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using Cathepsin D antibody (21327-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using Cathepsin D antibody (21327-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 21327-1-PBS in a different storage buffer formulation.