### For Research Use Only

# UHRF1 Polyclonal antibody

Catalog Number: 21402-1-AP

Featured Product

22 Publications



**Purification Method:** 

WB 1:500-1:2000

IF/ICC 1:50-1:500

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

Catalog Number:

GenBank Accession Number:

21402-1-AP

GeneID (NCBI):

BC113875

150ul, Concentration: 550 ug/ml by

29128

Nanodrop and 333 ug/ml by Bradford  $\,$  UNIPROT ID:

method using BSA as the standard;

Q96T88 Full Name:

Source: Rabbit

ubiquitin-like with PHD and ring

Isotype:

finger domains 1 Calculated MW:

Immunogen Catalog Number:

806 aa, 91 kDa Observed MW:

AG16111

91-100 kDa

**Applications** 

**Tested Applications:** 

WB, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, CoIP, RIP

Species Specificity:

human

**Cited Species:** 

human, mouse

Positive Controls:

WB: HuH-7 cells, HeLa cells, MCF-7 cells

IP: HeLa cells,

IF/ICC: HepG2 cells,

## **Background Information**

UHRF1, also named as ICBP90, NP95 and RNF106, is a putative E3 ubiquitin-protein ligase. It may participate in methylation-dependent transcriptional regulation. UHRF1 binds to inverted 5'-CCAAT-3' box 2 in the TOP2A promoter, and activates TOP2A expression. It is important for G1/S transition and may be involved in DNA repair and chromosomal stability. UHRF1's ability to repress its direct target gene expression is dependent on PHD(UHRF1) binding to unmodified H3R2. In addition, UHRF1 is a novel metabolic guardian restricting AMPK activity(PMID: 34907338).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Leyang Xiang	32808348	Cancer Sci	WB
Wei Deng	25833338	Cell Physiol Biochem	WB
Federica Morani	33800494	Int J Mol Sci	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 21402-1-AP (UHRF1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of UHRF1 antibody (21402-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-UHRF1 transfected HEK-293T



IP result of anti-UHRF1 (IP:21402-1-AP, 4ug; Detection:21402-1-AP 1:1000) with HeLa cells lysate 3200ug.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using UHRF1 antibody (21402-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).