

For Research Use Only

MRC2 Polyclonal antibody

Catalog Number: 21410-1-AP



Basic Information

Catalog Number: 21410-1-AP	GenBank Accession Number: BC146647	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 ug/ml by Nanodrop;	GeneID (NCBI): 9902	Recommended Dilutions: WB 1:1000-1:8000
Source: Rabbit	UNIPROT ID: Q9UBG0	
Isotype: IgG	Full Name: mannose receptor, C type 2	
Immunogen Catalog Number: AG15570	Calculated MW: 1479 aa, 167 kDa	
	Observed MW: 170 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, Saos-2 cells, SH-SY5Y cells
Species Specificity: human	

Background Information

C-type mannose receptor 2 (MRC2), also known as CD280, CLEC13E, ENDO180, KIAA0709, or UPARAP, is a type I transmembrane glycoprotein that belongs to the macrophage mannose receptor protein family (PMID: 10683150; 10636902; 9734811). It contains an N-terminal cysteine-rich domain, a fibronectin type II domain (FNII), eight C-type-lectin-like domains, and a transmembrane region followed by a small cytoplasmic domain (PMID: 34012764). MRC2 is a recycling endocytic receptor that functions in both these cellular activities by promoting cell migration and uptake of collagens for intracellular degradation (PMID: 17974964). MRC2 is expressed in several cells and organs including fibroblasts, endothelial cells, and macrophages. Altered MRC2 expression has been associated with the development and progression of different cancers, including head and neck squamous cell carcinoma, breast cancer, hepatocellular cancer, and prostate cancer (PMID: 36188226; 37666484; 17974964; 25162823).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21410-1-AP (MRC2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.