For Research Use Only

NRCAM Polyclonal antibody Catalog Number:21608-1-AP Featured Product 5

Featured Product 5 Publications




Basic Information	Catalog Number: 21608-1-AP	GenBank Accession Number: BC098401	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul, Concentration: 300 ug/ml by	4897	WB 1:500-1:2000
	Nanodrop and 233 ug/ml by Bradford method using BSA as the standard;	UNIPROT ID: Q92823	IHC 1:200-1:800 IF/ICC 1:50-1:500
	Source: Rabbit	Full Name: neuronal cell adhesion molecule Calculated MW:	
	Isotype:		
	IgG	1304 aa, 144 kDa	
	Immunogen Catalog Number: AG16237	Observed MW: 130-136 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, IF/ICC, ELISA Cited Applications:	WB: mouse brain ussue, numan brain ussue, nek-293	
	WB, IHC, IF		
	Species Specificity: human, mouse, rat	IF/ICC : HEK-293 cells,	
	Cited Species: human, mouse		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen	
	Neuronal cell adhesion molecule (NRCAM) is a member of the L1 subfamily of cell adhesion molecules (CAMs) that belong to the immunoglobulin superfamily (PMID: 11329126). NRCAM is a transmembrane protein composed of six Ig-like domains and five fibronectin type-III repeats in the extracellular region, with a highly conserved cytoplasmic tail. It is mainly expressed in the nervous system and is involved in neuron-neuron adhesion and promotes directional signaling during axonal cone growth. NRCAM is also expressed outside the nervous system. Altered expression of NRCAM has been associated with tumor progression in diverse organs (PMID: 22182708).		
Background Information	belong to the immunoglobulin super Ig-like domains and five fibronectin to cytoplasmic tail. It is mainly express promotes directional signaling during	family (PMID: 11329126). NRCA type-III repeats in the extracell ed in the nervous system and is g axonal cone growth. NRCAM i	M is a transmembrane protein composed of siv ular region, with a highly conserved s involved in neuron-neuron adhesion and s also expressed outside the nervous system.
	belong to the immunoglobulin super Ig-like domains and five fibronectin t cytoplasmic tail. It is mainly express promotes directional signaling during Altered expression of NRCAM has bee	family (PMID: 11329126). NRCA type-III repeats in the extracell ed in the nervous system and is g axonal cone growth. NRCAM i	M is a transmembrane protein composed of six ular region, with a highly conserved s involved in neuron-neuron adhesion and s also expressed outside the nervous system. ession in diverse organs (PMID: 22182708).
	belong to the immunoglobulin superf Ig-like domains and five fibronectin to cytoplasmic tail. It is mainly express promotes directional signaling during Altered expression of NRCAM has been Author Put	family (PMID: 11329126). NRCA type-III repeats in the extracell ed in the nervous system and is g axonal cone growth. NRCAM i en associated with tumor progre	M is a transmembrane protein composed of sizular region, with a highly conserved s involved in neuron-neuron adhesion and s also expressed outside the nervous system. ession in diverse organs (PMID: 22182708). Application
Background Information	belong to the immunoglobulin superf Ig-like domains and five fibronectin to cytoplasmic tail. It is mainly express promotes directional signaling during Altered expression of NRCAM has been Author Put Zhen-Huang Chen 329	family (PMID: 11329126). NRCA type-III repeats in the extracell ed in the nervous system and is g axonal cone growth. NRCAM i en associated with tumor progre med ID Journal	M is a transmembrane protein composed of sivular region, with a highly conserved sinvolved in neuron-neuron adhesion and s also expressed outside the nervous system. ession in diverse organs (PMID: 22182708).
	belong to the immunoglobulin superf Ig-like domains and five fibronectin to cytoplasmic tail. It is mainly express promotes directional signaling during Altered expression of NRCAM has been Author Put Zhen-Huang Chen 329 Michela Lupia 295	family (PMID: 11329126). NRCA type-III repeats in the extracell ed in the nervous system and is g axonal cone growth. NRCAM i en associated with tumor progre omed ID Journal 018999 Mol Cell Neur	M is a transmembrane protein composed of sivular region, with a highly conserved sinvolved in neuron-neuron adhesion and s also expressed outside the nervous system. ession in diverse organs (PMID: 22182708).
	belong to the immunoglobulin superf Ig-like domains and five fibronectin to cytoplasmic tail. It is mainly express promotes directional signaling during Altered expression of NRCAM has been Author Put Zhen-Huang Chen 329 Michela Lupia 295	family (PMID: 11329126). NRCA type-III repeats in the extracell ed in the nervous system and is g axonal cone growth. NRCAM i en associated with tumor progre omed ID Journal 018999 Mol Cell Neur 551673 Stem Cell Rep 150622 Front Oncol er shipment.	M is a transmembrane protein composed of sizular region, with a highly conserved sinvolved in neuron-neuron adhesion and s also expressed outside the nervous system. ession in diverse organs (PMID: 22182708).

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data







Various lysates were subjected to SDS PAGE followed by western blot with 21608-1-AP (NRCAM antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21608-1-AP (NRCAM antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21608-1-AP (NRCAM antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using 21608-1-AP (NRCAM antibody) at dilution of 1:50 and Alexa Fluor 488conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).