

AKT3 Polyclonal antibody

Catalog Number: 21641-1-AP

13 Publications

Basic Information

Catalog Number: 21641-1-AP	GenBank Accession Number: BC121154	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 760 µg/ml by Nanodrop;	GeneID (NCBI): 10000	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
Source: Rabbit	Full Name: v-akt murine thymoma viral oncogene homolog 3 (protein kinase B, gamma)	
Isotype: IgG	Calculated MW: 465 aa, 54 kDa	
Immunogen Catalog Number: AG16298	Observed MW: 60 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Cited Applications: IHC, WB	WB : mouse brain tissue, rat brain tissue
Species Specificity: human, mouse, rat	IHC : human tonsillitis tissue,
Cited Species: human	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

AKT3, also named as PKBG, is one of 3 closely related serine/threonine-protein kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which regulate many processes including metabolism, proliferation, cell survival, growth and angiogenesis. AKT3 is a key modulator of several tumors like melanoma, glioma and ovarian cancer. Active AKT3 increases progressively during melanoma tumor progression with highest levels present in advanced-stage metastatic melanomas.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Yang	31515938	J Cell Mol Med	WB
Yong-Xia Wang	30186040	Cancer Cell Int	WB
Wei Sang	27329362	Cell Immunol	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21641-1-AP (AKT3 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 21641-1-AP (AKT3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).