

For Research Use Only

NFAT5 Polyclonal antibody

Catalog Number: 21713-1-AP



Basic Information

Catalog Number: 21713-1-AP	GenBank Accession Number: BC146765	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 ug/ml by Nanodrop;	GeneID (NCBI): 10725	Recommended Dilutions: WB 1:1000-1:4000
Source: Rabbit	UNIPROT ID: O94916	IHC 1:50-1:500 IF/ICC 1:20-1:200
Isotype: IgG	Full Name: nuclear factor of activated T-cells 5, tonicity-responsive	
Immunogen Catalog Number: AG16341	Calculated MW: 1531 aa, 166 kDa	
	Observed MW: 200 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:
Species Specificity: human, mouse, rat, monkey	WB : HeLa cells, HepG2 cells, Jurkat cells, U2OS cells, COS-7 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse skeletal muscle tissue, human urothelial carcinoma tissue, human thyroid cancer tissue, mouse brain tissue, rat brain tissue
	IF/ICC : HepG2 cells,

Background Information

NFAT5 (Nuclear factor of activated T-cells 5) is the member of the Rel family of transcription factors, including NF- κ B and NFAT1-4, which plays central roles in inducible gene expression during the immune response (PMID: 21629436). NFAT5 also acts as a novel regulator of mammalian anti-pathogen responses, regulating TLR-induced gene expression in primary macrophages induced by Toll-like receptors (PMID: 22312110). NFAT5 is involved in the transcriptional regulation of osmoprotective and inflammatory genes (PMID: 10377394).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21713-1-AP (NFAT5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 21713-1-AP (NFAT5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 21713-1-AP (NFAT5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 21713-1-AP (NFAT5 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).