For Research Use Only

PI3 Kinase p110 Beta Polyclonal antibody

Catalog Number:21739-1-AP


17 Publications




Basic Information	Catalog Number: 21739-1-AP	GenBank Accession Number: BC114432	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 387 µg/ml by	5291	IHC 1:20-1:200
	Bradford method using BSA as the standard;	UNIPROT ID:	IF/ICC 1:50-1:500
	Source:	P42338 Full Name:	
	Rabbit	phosphoinositide-3-kinase, catalytic, beta polypeptide	
	Isotype:		
	lgG	Calculated MW:	
	Immunogen Catalog Number: AG16456	1070 aa, 123 kDa	
		Observed MW: 110 kDa	
Applications	Tested Applications:	Positive Controls:	
	IF, IHC, ELISA	Inc : numan tiver cancer tissue, numan	
	Cited Applications: WB, IHC, IF	IF/ICC : HeLa cells,	
	Species Specificity: human		
	Cited Species: human, mouse, rat		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
	PIK3CB(phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit beta isoform) is also named as PIK3C1, PI3K-beta, p110beta. The gene encodes a 1070 amino acid protein which belongs to the PI3/PI4-kinase family. Phosphoinositide 3-kinases (PI3Ks) have been implicated as participants in signaling pathways regulating cell growth by virtue of their activation in response to various mitogenic stimuli. The class I PI3 kinases are heterodimers composed of 110 kDa catalytic subunits that associate with regulatory adaptor proteins. Four class I catalytic subunits have been identified, PIK3CA (p110a), PIK3CB (p110β), PIK3CD (p110δ) and PIK3CG (p110γ) (PMID:19177002).		
Background Information	PI3K-beta, p110beta. The gene encod Phosphoinositide 3-kinases (PI3Ks) h growth by virtue of their activation ir heterodimers composed of 110 kDa c catalytic subunits have been identifi	es a 1070 amino acid protein wh ave been implicated as participa n response to various mitogenic s atalytic subunits that associate v	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are <i>i</i> th regulatory adaptor proteins. Four class
	PI3K-beta, p110beta. The gene encod Phosphoinositide 3-kinases (PI3Ks) h growth by virtue of their activation ir heterodimers composed of 110 kDa c catalytic subunits have been identifi (PMID:19177002).	es a 1070 amino acid protein wh ave been implicated as participa n response to various mitogenic s atalytic subunits that associate v ed, PIK3CA (p110a), PIK3CB (p110	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are rith regulatory adaptor proteins. Four class ββ), PIK3CD (p110δ) and PIK3CG (p110γ)
	PI3K-beta, p110beta. The gene encod Phosphoinositide 3-kinases (PI3Ks) h growth by virtue of their activation in heterodimers composed of 110 kDa c catalytic subunits have been identifi (PMID:19177002).	es a 1070 amino acid protein wh ave been implicated as participa n response to various mitogenic s atalytic subunits that associate v ed, PIK3CA (p110a), PIK3CB (p110 pmed ID Journal	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are <i>i</i> th regulatory adaptor proteins. Four class
Background Information	PI3K-beta, p110beta. The gene encod Phosphoinositide 3-kinases (PI3Ks) h growth by virtue of their activation in heterodimers composed of 110 kDa c catalytic subunits have been identifi (PMID:19177002). Author Put Zhaomeng Wu 296	es a 1070 amino acid protein wh ave been implicated as participa or response to various mitogenic s atalytic subunits that associate v ed, PIK3CA (p110a), PIK3CB (p110 pmed ID Journal 520175 Mol Med Rep	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are <i>i</i> th regulatory adaptor proteins. Four class ββ, PIK3CD (p110δ) and PIK3CG (p110γ) Application
	PI3K-beta, p110beta. The gene encode Phosphoinositide 3-kinases (PI3Ks) he growth by virtue of their activation in heterodimers composed of 110 kDa c catalytic subunits have been identified (PMID:19177002). Author Put Zhaomeng Wu 296 Yuan-Yuan Gao 350	es a 1070 amino acid protein wh ave been implicated as participa nesponse to various mitogenic s atalytic subunits that associate v ed, PIK3CA (p110a), PIK3CB (p110 pmed ID Journal 520175 Mol Med Rep 047352 Int J Ophthalmo	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are <i>i</i> th regulatory adaptor proteins. Four class oβ), PIK3CD (p110δ) and PIK3CG (p110γ) Application
	PI3K-beta, p110beta. The gene encode Phosphoinositide 3-kinases (PI3Ks) he growth by virtue of their activation in heterodimers composed of 110 kDa c catalytic subunits have been identified (PMID:19177002). Author Put Zhaomeng Wu 296 Yuan-Yuan Gao 350	es a 1070 amino acid protein wh ave been implicated as participa or response to various mitogenic s atalytic subunits that associate v ed, PIK3CA (p110a), PIK3CB (p110 pmed ID Journal 520175 Mol Med Rep	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are <i>i</i> th regulatory adaptor proteins. Four class oβ), PIK3CD (p110δ) and PIK3CG (p110γ) Application
	PI3K-beta, p110beta. The gene encode Phosphoinositide 3-kinases (PI3Ks) he growth by virtue of their activation in heterodimers composed of 110 kDa c catalytic subunits have been identified (PMID:19177002). Author Put Zhaomeng Wu 296 Yuan-Yuan Gao 350	es a 1070 amino acid protein wh ave been implicated as participa or response to various mitogenic s atalytic subunits that associate v ed, PIK3CA (p110a), PIK3CB (p110 ormed ID Journal 520175 Mol Med Rep 047352 Int J Ophthalmo 343164 J Zhejiang Univ er shipment. % glycerol pH 7.3.	ch belongs to the PI3/PI4-kinase family. nts in signaling pathways regulating cell timuli. The class I PI3 kinases are <i>i</i> th regulatory adaptor proteins. Four class oβ), PIK3CD (p110δ) and PIK3CG (p110γ) Application

T: 1 (1888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human liver cancer using 21739-1-AP (PI3K p110(beta) antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver cancer using 21739-1-AP (PI3K p110(beta) antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 21739-1-AP (PI3K p110(beta) antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).