For Research Use Only

VNN1 Polyclonal antibody

Catalog Number: 21745-1-AP

Featured Product

12 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

21745-1-AP

Size:

Source:

Rabbit

Isotype:

BC096268 GeneID (NCBI):

8876

Nanodrop and 333 ug/ml by Bradford $\,$ UNIPROT ID:

095497

Full Name:

vanin 1 Calculated MW:

513 aa, 57 kDa Immunogen Catalog Number:

150ul, Concentration: 333 ug/ml by

method using BSA as the standard;

Observed MW: AG16501

55-72 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC, IF, CoIP

Species Specificity:

human, mouse **Cited Species:**

human, mouse

Positive Controls:

WB: rat kidney tissue, BxPC-3 cells, mouse kidney

tissue, PC-13 cells

Background Information

VNN1(Vascular non-inflammatory molecule 1) is also named as Vanin-1, Pantetheinase, Tiff66. It is a GPI-anchored 70-kDa protein with a high degree of homology with GPI-80 (also known as VNN2), a molecule expressed by human phagocytes and involved in leukocyte adhesion and migration. Vanin-1 is predominantly expressed by resident tissue cells and acts at the level of epithelial cells and surrounding hematopoietic cells by paracrine 2- $Mercaptoethylamine\ release (PMID:17163446).\ The\ full\ length\ VNN1\ has\ a\ signal\ peptide,\ a\ propeptide\ and\ six$ glycosylation sites. According to some authors, the variations which varied between 55 and 72kDa may be due to different degrees of glycosylation of the protein and cleavage (JOSE A et al. 2001).

Notable Publications

Author	Pubmed ID	Journal	Application
Povero Davide D	24106341	Sci Signal	WB
Hong Xiang	33754072	Theranostics	ColP
Yanxiang Wu	35234362	Proteomics	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21745-1-AP (VNN1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.

BxPC-3 cells were subjected to SDS PAGE followed by western blot with 21745-1-AP (VNN1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.