

DELE1 Polyclonal antibody

Catalog Number: 21904-1-AP

2 Publications

Basic Information

Catalog Number: 21904-1-AP	GenBank Accession Number: BC007855	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop and 360 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9812	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	Full Name: KIAA0141	
Isotype: IgG	Calculated MW: 515 aa, 56 kDa	
Immunogen Catalog Number: AG14615	Observed MW: 56 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse lung tissue,
Cited Applications: WB	
Species Specificity: human, mouse	
Cited Species: mouse	

Background Information

DELE1, also known as KIAA0141, is a 515 amino acid protein, which localizes in Mitochondrion and is detected in liver, skeletal muscle, kidney, pancreas, spleen, thyroid, testis, ovary, small intestine and colon. DELE1 is essential for the induction of death receptor-mediated apoptosis through the regulation of caspase activation. Mitochondrial malfunction needs to be relayed to the cytosol, where an integrated stress response is triggered by the phosphorylation of eukaryotic translation initiation factor 2α (eIF2α) in mammalian cells. eIF2α phosphorylation is mediated by the four eIF2α kinases GCN2, HRI, PERK and PKR, which are activated by diverse types of cellular stress. OMA1, a mitochondrial stress-activated protease; and DELE1, a little-characterized protein that we found was associated with the inner mitochondrial membrane. Mitochondrial stress stimulates OMA1-dependent cleavage of DELE1 and leads to the accumulation of DELE1 in the cytosol, where it interacts with HRI and activates the eIF2α kinase activity of HRI. In addition, DELE1 is required for ATF4 translation downstream of eIF2α phosphorylation.

Notable Publications

Author	Pubmed ID	Journal	Application
Jiahui Tang	36290724	Antioxidants (Basel)	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse lung tissue were subjected to SDS PAGE followed by western blot with 21904-1-AP (KIAA0141 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.