

For Research Use Only

ACC1 Polyclonal antibody

Catalog Number: 21923-1-AP

Featured Product

170 Publications



Basic Information

Catalog Number:

21923-1-AP

Size:

150ul, Concentration: 800 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG16452

GenBank Accession Number:

BC137287

GeneID (NCBI):

31

UNIPROT ID:

Q13085

Full Name:

acetyl-Coenzyme A carboxylase alpha

Calculated MW:

2383 aa, 275 kDa

Observed MW:

250 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:1000-1:20000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

IF/ICC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, RIP, IHC-P

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, pig, chicken, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, HepG2 cells, mouse brain tissue, rat brain tissue

IP: HepG2 cells,

IHC: mouse skeletal muscle tissue, mouse brain tissue

IF/ICC: HeLa cells,

Background Information

ACACA (Acetyl-CoA carboxylase 1, ACC), also named as ACAC, ACC1 and ACCA, belongs to the biotin containing enzyme family. It catalyzes the synthesis of malonyl-CoA, which is an intermediate substrate playing a pivotal role in the regulation of fatty acid metabolism and energy production. ACACA is involved in the biosynthesis of fatty acids, and malonyl-CoA produced is used as a building block to extend the chain length of fatty acids by fatty acid synthase (FAS) (PMID:19900410). It has 4 isoforms produced by alternative promoter usage with the molecular weight between 260 kDa and 270 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Lihua Luo	34593005	J Nanobiotechnology	WB
Shifeng Pan	29152131	Oncotarget	WB
Zhongwen Feng	33182043	Int Immunopharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 21923-1-AP (ACC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using ACC1 antibody (21923-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




WB result of ACC1 antibody (21923-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACC1 transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 21923-1-AP (ACC1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-ACC1 (IP:21923-1-AP, 4ug; Detection:21923-1-AP 1:2000) with HepG2 cells lysate 1800 ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21923-1-AP (ACC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).