

For Research Use Only

# FTCD Polyclonal antibody

Catalog Number: 21959-1-AP

4 Publications



## Basic Information

<b>Catalog Number:</b> 21959-1-AP	<b>GenBank Accession Number:</b> BC136395	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 500 µg/ml by Nanodrop and 480 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 10841	<b>Recommended Dilutions:</b> WB 1:4000-1:16000 IHC 1:250-1:1000
<b>Source:</b> Rabbit	<b>Full Name:</b> formiminotransferase cyclodeaminase	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 541 aa, 59 kDa	
<b>Immunogen Catalog Number:</b> AG16646	<b>Observed MW:</b> 53-62 kDa	

## Applications

<b>Tested Applications:</b> IHC, WB, ELISA	<b>Positive Controls:</b>
<b>Cited Applications:</b> IF, IHC, WB	<b>WB :</b> mouse liver tissue, rat liver tissue
<b>Species Specificity:</b> human, mouse, rat	<b>IHC :</b> human liver tissue,
<b>Cited Species:</b> human	

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
Shizhe Yu	35813194	Front Cell Dev Biol	IF
Jiajia Chen	31203308	Med Sci Monit	WB,IHC
Huimin Shen	33380233	Bioengineered	IHC

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21959-1-AP (FTCD antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 21959-1-AP (FTCD antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 21959-1-AP (FTCD antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).