For Research Use Only

DULLARD Polyclonal antibody

Catalog Number:22088-1-AP



Basic Information	Catalog Number: 22088-1-AP	GenBank Accession Number BC009295	er: Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 600 ug/ml by Nanodrop and 327 ug/ml by Bradford method using BSA as the standard;	23399 UNIPROT ID: 095476	WB 1:200-1:1000 IHC 1:50-1:500 IF/ICC 1:50-1:500
	Source: Rabbit	Full Name: dullard homolog (Xenopus laevis)	
	Isotype:Calculated MW:IgG244 aa, 28 kDa		
	Immunogen Catalog Number: AG17288	Observed MW: 28 kDa	
Applications	Tested Applications: WB, IHC, IF/ICC, EUSA	Pos	itive Controls:
	Species Specificity: human, mouse		: PC-3 cells, mouse skeletal muscle tissue : human ovary cancer tissue,
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		CC : PC-3 cells,
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	0, 1	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 22088-1-AP (DULLARD antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using DULLARD antibody (22088-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).

PC-3 cells were subjected to SDS PAGE followed by western blot with 22088-1-AP (DULLARD Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.