For Research Use Only

TIGAR Polyclonal antibody

Catalog Number:22136-1-AP

6 Publications



IHC: mouse colon tissue, mouse small intestine tissue

Basic Information

Catalog Number: GenBank Accession Number: 22136-1-AP

Purification Method: Antigen affinity purification BC012340 GeneID (NCBI): Recommended Dilutions:

150ul, Concentration: 845 µg/ml by 57103 Nanodrop and 845 µg/ml by Bradford Full Name:

WB 1:500-1:2000 IHC 1:50-1:500

method using BSA as the standard;

chromosome 12 open reading frame 5

Calculated MW: Rabbit 270 aa, 30 kDa Isotype: Observed MW: IgG 30 kDa

Immunogen Catalog Number:

AG17532

WB

Applications

Tested Applications: Positive Controls: IHC, WB, ELISA

WB: mouse skeletal muscle tissue, rat skeletal muscle **Cited Applications:**

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoming Chai	36399798	Biochem Biophys Res Commun	WB
Shan Liu	35165280	Cell Death Dis	WB
Yogita Kanan	35134532	Free Radic Biol Med	WB

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 22136-1-AP (TIGAR antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 22136-1-AP (TIGAR antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 22136-1-AP (TIGAR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.