#### For Research Use Only

# iNOS Polyclonal antibody

Catalog Number:22226-1-AP 150 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

22226-1-AP

GeneID (NCBI):

Antigen affinity purification

150ul , Concentration: 600 ug/ml by

Recommended Dilutions: WB 1:200-1:1000

Nanodrop:

**UNIPROT ID:** P35228

BC130283

Rabbit Full Name:

nitric oxide synthase 2, inducible

Isotype: IgG

Calculated MW:

Immunogen Catalog Number:

1153 aa, 131 kDa

AG17567

Observed MW:

110-130 kDa, 65-70 kDa

**Applications** 

**Tested Applications:** WB, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse **Cited Species:** 

human, mouse, rat, chicken

Positive Controls:

WB: RAW 264.7 cells,

## **Background Information**

NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70kD band can be detected by western (PMID:11786228).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Mao Kobayashi	32992650	Int J Mol Sci	WB
Jingming Zhuang	36161709	Andrology	WB
Dewei He	36139502	Cells	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

### **Selected Validation Data**



Untreated and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 22226-1-AP (NOS2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



1X10^6 LPS treated RAW 264.7 cells were intracellularly stained with 0.4 ug Anti-Human iNOS (22226-1-AP) and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).