

For Research Use Only

iNOS Polyclonal antibody

Catalog Number: 22226-1-AP **18 Publications**



Basic Information

Catalog Number: 22226-1-AP	GenBank Accession Number: BC130283	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 1000 µg/ml by Nanodrop and 533 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 4843	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	Full Name: nitric oxide synthase 2, inducible	
Isotype: IgG	Calculated MW: 1153 aa, 131 kDa	
Immunogen Catalog Number: AG17567	Observed MW: 110-130 kDa, 65-70 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : RAW 264.7 cells,
Cited Applications: IF, IHC, WB	
Species Specificity: human, mouse	
Cited Species: human, mouse, rat	

Background Information

NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70kD band can be detected by western (PMID:11786228).

Notable Publications

Author	Pubmed ID	Journal	Application
Mao Kobayashi	32992650	Int J Mol Sci	WB
Qicai Liu	33067662	Inflammation	IF
Ziru Zhang	31790900	Phytomedicine	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Untreated and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 22226-1-AP (NOS2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.