For Research Use Only

CCL24/Eotaxin 2 Polyclonal antibody

Catalog Number:22306-1-AP 2 Publications



Basic Information	Catalog Number: 22306-1-AP	GenBank Accession Number: NM_002991	Purification Method: Antigen Affinity purified
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 500 ug/ml by Nanodrop and 340 ug/ml by Bradford method using BSA as the standard;	6369	WB 1:200-1:1000
		UNIPROT ID: 000175	IHC 1:50-1:500
	Source: Rabbit	Full Name: chemokine (C-C motif) ligand 24	
	lsotype: IgG	Calculated MW: 13 kDa	
		Observed MW: 13 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, ELISA Cited Applications:	WB : mouse thymus tissue,	
	WB, IF, IHC	IHC : human breast cancer tissue,	
	Species Specificity: human, mouse		
	Cited Species: human, mouse		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
	Chemokine CCL24 is the second member of eotaxins, a group of eosinophils' selectively chemoattractants. CCL24 i constitutively expressed in fibroblasts, epithelial cells, macrophages. CCL24 has only one receptor CCR3, which mainly presents on eosinophils and was also found on basophils, monocytes, Th2 lymphocytes, epithelial cells and airway smooth muscles. CCL24 displays chemotactic activity on resting T lymphocytes, a minimal activity on neutrophils, and is negative on monocytes and activated T lymphocytes.		
Background Information	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa	ts, epithelial cells, macrophages. vas also found on basophils, mon ays chemotactic activity on restin	CCL24 has only one receptor CCR3, which ocytes, Th2 lymphocytes, epithelial cells an g T lymphocytes, a minimal activity on
	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa neutrophils, and is negative on mono	ts, epithelial cells, macrophages. vas also found on basophils, mon ays chemotactic activity on restin	CCL24 has only one receptor CCR3, which ocytes, Th2 lymphocytes, epithelial cells an g T lymphocytes, a minimal activity on
Background Information	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa neutrophils, and is negative on mono Author Pub	ts, epithelial cells, macrophages. vas also found on basophils, mon ays chemotactic activity on restin cytes and activated T lymphocyte	CCL24 has only one receptor CCR3, which bcytes, Th2 lymphocytes, epithelial cells an g T lymphocytes, a minimal activity on is.
	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa neutrophils, and is negative on mono Author Pub Youdi Wang 342	ts, epithelial cells, macrophages. vas also found on basophils, mon- ays chemotactic activity on restin cytes and activated T lymphocyte med ID Journal	CCL24 has only one receptor CCR3, which bcytes, Th2 lymphocytes, epithelial cells an g T lymphocytes, a minimal activity on es. Application
	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa neutrophils, and is negative on mono Author Pub Youdi Wang 342	ts, epithelial cells, macrophages. vas also found on basophils, mon ays chemotactic activity on restin cytes and activated T lymphocyte med ID Journal 21188 Dis Markers	CCL24 has only one receptor CCR3, which boytes, Th2 lymphocytes, epithelial cells ar g T lymphocytes, a minimal activity on es. Application WB
Notable Publications	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa neutrophils, and is negative on mono Author Pub Youdi Wang 342 Jiahao He 382 Storage: Stora at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°	ts, epithelial cells, macrophages. vas also found on basophils, mon ays chemotactic activity on restin cytes and activated T lymphocyte med ID Journal 21188 Dis Markers 67432 Cell Death Dis er shipment. % glycerol pH 7.3.	CCL24 has only one receptor CCR3, which boytes, Th2 lymphocytes, epithelial cells ar g T lymphocytes, a minimal activity on es. Application WB
	constitutively expressed in fibroblast mainly presents on eosinophils and w airway smooth muscles. CCL24 displa neutrophils, and is negative on mono Author Pub Youdi Wang 342 Jiahao He 382 Storage: Stora at -20°C. Stable for one year after Storage Buffer:	ts, epithelial cells, macrophages. vas also found on basophils, mon ays chemotactic activity on restin cytes and activated T lymphocyte med ID Journal 21188 Dis Markers 67432 Cell Death Dis er shipment. % glycerol pH 7.3.	CCL24 has only one receptor CCR3, which boytes, Th2 lymphocytes, epithelial cells an g T lymphocytes, a minimal activity on es. Application WB

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse thymus tissue were subjected to SDS PAGE followed by western blot with 22306-1-AP (CCL24/Eotaxin 2 antibody at dilution of 1:300 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 22306-1-AP (CCL24/Eotaxin 2 antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 22306-1-AP (CCL24/Eotaxin 2 antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).