

For Research Use Only

BTG2 Polyclonal antibody

Catalog Number: 22339-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number: 22339-1-AP	GenBank Accession Number: BC105949	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 350 µg/ml by Nanodrop and 260 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 7832	Recommended Dilutions: IHC 1:50-1:500 IF 1:200-1:800
Source: Rabbit	Full Name: BTG family, member 2	
Isotype: IgG	Calculated MW: 158 aa, 17 kDa	
Immunogen Catalog Number: AG17819		

Applications

Tested Applications: FC, IF, IHC, ELISA	Positive Controls:
Cited Applications: IF, IHC, WB	IHC: mouse small intestine tissue, human kidney tissue
Species Specificity: human, mouse	IF: Neuro-2a cells,
Cited Species: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Protein BTG2, the anti-proliferative human homolog of murine Pc3/Tis21, belongs to the BTG/Tob family. This family has structurally related proteins that appear to have anti-proliferative properties. It has been demonstrated that BTG2 is p53-inducible and is involved in cell cycle control and cellular response to DNA damage (PMID: 8944033). As a transcriptional co-regulator, BTG2 has been shown to interact with CAF1 and POP2, and works as a co-activator of ERα-mediated transcription via a CCR4-like complex (PMID: 18974182).

Notable Publications

Author	Pubmed ID	Journal	Application
Shi Li	31885582	J Oncol	WB
Chen-Song Huang	29098552	Dig Dis Sci	IHC
Aitao Nai	35372503	Front Mol Biosci	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse small intestine tissue slide using 22339-1-AP (BTG2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse small intestine tissue slide using 22339-1-AP (BTG2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using BTG2 antibody (22339-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human BTG2 (22339-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).