## For Research Use Only

## CD79a Polyclonal antibody

Catalog Number:22349-1-AP 1 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC113733

**Purification Method:** Antigen affinity purification

22349-1-AP Size:

GeneID (NCBI):

150ul , Concentration: 600 ug/ml by

Recommended Dilutions: WB 1:500-1:2000

Nanodrop:

**UNIPROT ID:** P11912

IHC 1:200-1:800 IF-P 1:50-1:500

Rabbit

IgG

Full Name:

Isotype:

CD79a molecule, immunoglobulin-

associated alpha

Immunogen Catalog Number:

Calculated MW:

AG17924

226 aa. 25 kDa

Observed MW:

38-42 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF-P, ELISA

**Cited Applications:** 

Species Specificity:

human

Cited Species: mouse

TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

Note-IHC: suggested antigen retrieval with

buffer pH 6.0

Positive Controls:

WB: Daudi cells, Raji cells, Ramos cells

IHC: human tonsillitis tissue, human lymphoma tissue

IF-P: human tonsillitis tissue,

**Background Information** 

CD79a, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79A is expressed in B cell and localizes in the cell membrane. CD79A is required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the Bcell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79A is also required for BCR surface expression and for efficient differentiation of proand pre-B-cells.

**Notable Publications** 

Author **Pubmed ID** Journal Application Zulikeyan Manafu 38723832 Int J Biol Macromol IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 22349-1-AP (CD79A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 22349-1-AP (CD79A antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 22349-1-AP (CD79A antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD79A antibody (22349-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD79A antibody (22349-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).