

For Research Use Only

WDR33 Polyclonal antibody

Catalog Number: 22614-1-AP

1 Publications



Basic Information

Catalog Number: 22614-1-AP	GenBank Accession Number: BC010283	Purification Method: Antigen Affinity purified
Size: 150ul , Concentration: 1300 ug/ml by Nanodrop and 513 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 55339	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Source: Rabbit	UNIPROT ID: Q9C0J8	
Isotype: IgG	Full Name: WD repeat domain 33	
Immunogen Catalog Number: AG18382	Calculated MW: 1336 aa, 146 kDa	
	Observed MW: 170 kDa	

Applications

Tested Applications: WB, IP, ELISA	Positive Controls: WB : HepG2 cells, HeLa cells IP : HeLa cells,
Cited Applications: WB	
Species Specificity: human	
Cited Species: human	

Background Information

WD repeat domain 33 (WDR33), also named WDC146, is a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-aspartate (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This gene is highly expressed in testis and the protein is localized to the nucleus. This gene may play important roles in the mechanisms of cytodifferentiation and/or DNA recombination. This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-aspartate (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This gene is highly expressed in testis and the protein is localized to the nucleus. This gene may play important roles in the mechanisms of cytodifferentiation and/or DNA recombination. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. This antibody recognizes the full length WDR33 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Qixuan Lu	39190270	Front Med	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 22614-1-AP (WDR33 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

IP result of anti-WDR33 (IP:22614-1-AP, 4ug; Detection:22614-1-AP 1:1000) with HeLa cells lysate 3200ug.