

For Research Use Only

NOP10 Polyclonal antibody

Catalog Number: 22769-1-AP



Basic Information

Catalog Number: 22769-1-AP	GenBank Accession Number: BC063023	Purification Method: Antigen Affinity purified
Size: 150ul , Concentration: 1300 ug/ml by Nanodrop;	GeneID (NCBI): 55505	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:200-1:800
Source: Rabbit	UNIPROT ID: Q9NPE3	
Isotype: IgG	Full Name: NOP10 ribonucleoprotein homolog (yeast)	
Immunogen Catalog Number: AG18718	Calculated MW: 64 aa, 77 kDa	
	Observed MW: 10-12 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Species Specificity: human	WB : COLO 320 cells, HeLa cells, Jurkat cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : human pancreas cancer tissue,

Background Information

Nucleolar protein 10 (NOP10) is a component of the H/ACA small nucleolar ribonucleoprotein (snoRNP) complex and is essential for maintaining telomerase activity and RNA pseudouridylation. NOP10 is abnormally expressed in a variety of cancers such as breast cancer, pancreatic cancer, lung cancer, etc., and high expression is usually associated with poor prognosis(PMID: 33161513; 39538226).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 22769-1-AP (NOP10 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 22769-1-AP (NOP10 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 22769-1-AP (NOP10 antibody) at dilution of 1:400 (under 0x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).