For Research Use Only

Dectin-1 Polyclonal antibody

Catalog Number:22809-1-AP



Basic Information	Catalog Number: 22809-1-AP	GenBank Accession Number: BC013385	Purification Method: Antigen affinity purification		
	Size: 150ul , Concentration: 1000 ug/ml by Nanodrop and 547 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG18924	GenelD (NCBI): 64581	Recommended Dilutions: WB 1:500-1:1000		
		UNIPROT ID: Q9BXN2			
		Full Name: C-type lectin domain family 7, member A Calculated MW: 247 aa, 28 kDa			
				Observed MW: 45 kDa	
				Applications	Tested Applications: WB, ELISA Species Specificity: human
		WB : HL-60 cells, THP-1 cells, U-937 cells			
Background Information	Dectin-1, also known as CLEC7A or CD369, is a type II membrane receptor consisting of an extracellular C-terminal C-type lectin domain, a short stalk region, a single transmembrane domain and a short N-terminal cytoplasmic tail, which contains an immunoreceptor tyrosine-based activation motif (PMID: 10779524; 21612412). Dectin-1 is expressed on monocytes, macrophages, neutrophils, dendritic cells and a subpopulation of T cells (PMID: 12244185) Dectin-1 plays a role in the innate immune response. It functions as a pattern-recognition receptor and recognizes a variety of β -glucans from fungi, plants and bacteria. Dectin-1 may also participate in orchestrating the adaptive immune response (PMID: 21612412). The apparent molecular weight is larger than the calculated molecular weight due to glycosylation (PMID: 10779524).				
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°				
	Aliquoting is unnecessary for -20°C st	orage			

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 22809-1-AP (Dectin-1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.