

For Research Use Only

IL-4|1 Polyclonal antibody

Catalog Number: 23048-1-AP



Basic Information

Catalog Number: 23048-1-AP	GenBank Accession Number: BC131625	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 259307	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	Full Name: interleukin 4 induced 1	
Isotype: IgG	Calculated MW: 567 aa, 63 kDa	
Immunogen Catalog Number: AG19136	Observed MW: 70 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, Jurkat cells, mouse liver tissue
Species Specificity: human, mouse	

Background Information

Interleukin-4 Induced Gene 1 (IL4I1) is a secreted L-phenylalanine oxidase mainly produced by inflammatory antigen-presenting cells-in particular, macrophages present in T helper type 1 granulomas and in various types of tumors. In the last decade, it has been shown that IL4I1 is involved in the fine control of B- and T-cell adaptive immune responses. Preclinical models have revealed its role in cancer immune evasion. Recent clinical data highlight IL4I1 as a new potential prognostic marker in human melanoma. As a secreted enzyme, IL4I1 may represent an easily targetable molecule for cancer immunotherapy. IL4I1 protein has an estimated molecular weight of 75 kDa. (PMID: 31330829, PMID: 21469114)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 23048-1-AP (IL-41 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.