For Research Use Only

NCOA7 Polyclonal antibody

Catalog Number:23092-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number:

23092-1-AP

Size:

Source:

GenBank Accession Number:

BC137094

GeneID (NCBI):

135112

150ul , Concentration: 500 ug/ml by Nanodrop and 367 ug/ml by Bradford $\,$ UNIPROT ID:

method using BSA as the standard;

Q8NI08 Full Name:

Rabbit

nuclear receptor coactivator 7

Isotype: Calculated MW:

972 aa, 106 kDa

Immunogen Catalog Number:

Observed MW:

AG19315

120-130 kDa

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IF, IHC, CoIP

Species Specificity:

human

Cited Species:

human, mouse

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Positive Controls:

WB: HEK-293 cells, MCF-7 cells

IP: HEK-293 cells,

Background Information

 $Nuclear\ receptors\ for\ steroids, thy roid\ hormones\ and\ retinoic\ acids\ are\ ligand-dependent\ transcription\ factors\ that$ activate transcription through specific DNA binding sites in their target genes. NCoA-7 (nuclear receptor coactivator 7), also known as ESNA1 or ERAP140, is a 942 amino acid nuclear protein that enhances nuclear receptor transcriptional activities and coactivates several nuclear receptors including PPARG, ESR1, THRB and RARA. Highly expressed in brain and weakly expressed in pancreas, bladder, ovary, spinal cord, prostate, mammary gland, ovary, uterus and stomach, NCoA-7 is a member of the OXR1 family and contains one LysM repeat and a TLD domain. This antibody is a rabbit polyclonal antibody raised against the N-terminal of human NCOA7 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Maria Merkulova	29384414	Am J Physiol Renal Physiol	
Enrico Castroflorio	33340069	Cell Mol Life Sci	WB,CoIP
Jiayu Guo	37511343	Int J Mol Sci	WB,IHC,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 23092-1-AP (NCOA7 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-NCOA7 (IP:23092-1-AP, 4ug; Detection:23092-1-AP 1:1000) with HEK-293 cells lysate 3600ug.