### For Research Use Only

# MYD88 Polyclonal antibody

Catalog Number:23230-1-AP

Featured Product

212 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

23230-1-AP BC013589
Size: GeneID (NCBI):

150ul, Concentration: 600 ug/ml by 4615

Nanodrop; UNIPROT ID:
Source: Q99836
Rabbit Full Name:

Isotype: myeloid differentiation primary

IgG response gene (88)

Immunogen Catalog Number: Calculated MW:

AG19770 33 kDa

Observed MW:

33 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

# **Applications**

Tested Applications: WB, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity:

human
Cited Species:

human, pig, canine, monkey, chicken, bovine, sheep

#### Positive Controls:

WB: Jurkat cells, A549 cells, HepG2 cells, MCF-7 cells,

K-562 cells, Raji cells
IP: K-562 cells, A549 cells

# **Background Information**

MYD88 is a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. MYD88 is the key adaptor protein for the interleukin-1 receptor (IL-1R) and most Toll-like receptors (TLRs), which are essential for innate immunity and pathogen-associated molecular pattern recognition. Mutations in the MyD88 gene lead to the development of cancer in humans and mice suggesting that MyD88 also plays a cell autonomous role in tissue homeostasis.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Chang Liu	36230117	Foods	WB
Yangming Zhang	36191372	Biochem Biophys Res Commun	WB
Zhaohai Wen	36238295	Front Immunol	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 23230-1-AP (MYD88 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




WB result of MYD88 antibody (23230-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MYD88 transfected HepG2 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 23230-1-AP (MYD88 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-MYD88 (IP:23230-1-AP, 4ug; Detection:23230-1-AP 1:10000) with K-562 cells lysate 1360 ug.



IP result of anti-MYD88 (IP:23230-1-AP, 4ug; Detection:23230-1-AP 1:10000) with A549 cells lysate 640 ug.