For Research Use Only

PCF11 Polyclonal antibody

Catalog Number:23540-1-AP

in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

Featured Product

5 Publications




Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 23540-1-AP BC146778 Antigen Affinity purified GenelD (NCBI): Recommended Dilutions: Size: 150ul , Concentration: 600 ug/ml by 51585 WB 1:500-1:2000 Nanodrop: IP 0.5-4.0 ug for 1.0-3.0 mg of total UNIPROT ID: protein lysate Source 094913 IHC 1:200-1:800 Rabbit Full Name: Isotype PCF11, cleavage and polyadenylation factor subunit, homolog (S. lgG cerevisiae) Immunogen Catalog Number: Calculated MW: AG20263 1555 aa, 173 kDa Observed MW: 173 kDa **Positive Controls: Applications Tested Applications:** WB, IHC, IP, ELISA WB: K-562 cells, HeLa cells **Cited Applications:** IP : K-562 cells, WB. ColP IHC : human intrahepatic cholangiocarcinoma tissue, **Species Specificity:** human **Cited Species:** human, mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Background Information** In Saccharomyces cerevisiae, the cleavage/polyadenylation factor Pcf11 is a crucial regulatory factor required for recruiting polyadenylation machinery to elongating RNA polymerase II (RNAPII), and is necessary for correct transcriptional termination. Pcf11 (PCF11, cleavage and polyadenylation factor subunit, homolog (S. cerevisiae)), is a 1,555 amino acid nuclear protein that is a component of pre-mRNA cleavage complex II. It is suggested that Pcf11 is capable of promoting the dissociation of Pol II elongation complexes from DNA. Pcf11 contains a CTD-interaction domain that binds in a phospho-dependent manner to the heptad repeats within the RNA polymerase II CTD. The gene encoding Pcf11 is located on human chromosoem 11, which houses over 1,400 genes and comprises nearly 4% of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that maps to chromosome 11. This antibody is specific to the 173kd human PCF11 protein. **Notable Publications** Author Pubmed ID Journal Application Jihae Shin 34244761 Nucleic Acids Res WB Geneva R LaForce 35139363 Neuron ColP Biochimie 27060432 Asya Khleborodova WB Storage Storage Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage *** 20ul sizes contain 0.1% BSA For technical support and original validation data for this product please contact: This product is exclusively available under Proteintech T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com Group brand and is not available to purchase from any

other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human intrahepatic cholangiocarcinoma tissue slide using 23540-1-AP (PCF11 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

IP result of anti-PCF11 (IP:23540-1-AP, 4ug; Detection:23540-1-AP 1:1000) with K-562 cells lysate 2400 ug.

K-562 cells were subjected to SDS PAGE followed by western blot with 23540-1-AP (PCF 11 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.