## For Research Use Only

## FNDC5 Polyclonal antibody

Catalog Number:23995-1-AP

25 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

BC062297 Antigen affinity purification

GenelD (NCBI): Recommended Dilutions:

tration: 350 ug/ml by 252995 WB 1:1000-1:4000

150ul , Concentration: 350 µg/ml by 252995 WB 1:1000-1:400 Nanodrop; Full Name: IHC 1:200-1:800

Source: fibronectin type III domain containing

Rabbit 5

Isotype: Calculated MW:
IgG 212 aa, 24 kDa
Immunogen Catalog Number: Observed MW:
AG21195 25-30 kDa

**Applications** 

Tested Applications:

IHC, WB, ELISA

23995-1-AP

Cited Applications: IF, IHC, WB

Species Specificity: human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse liver tissue, mouse stomach tissue, mouse

**Purification Method:** 

skeletal muscle tissue, rat liver tissue IHC: mouse skeletal muscle tissue.

**Background Information** 

fibronectin type III domain containing 5(FNDC5)encodes 23kDa protein, which is a membrane protein that is cleaved and secreted as a newly identified hormone, irisin. The exercise induces muscle FNDC5 expression. Exercise-induced FNDC5 gene expression in muscles was accompanied by a parallel increase in the concentration of circulating irisin, which in turn activates adipocyte thermogenic programs, leading to mitochondrial heat production and energy expenditure.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Xianshengjie Lang	32987101	Brain Res Bull	WB
Wuyang He	32933582	Skelet Muscle	WB
Yi Yan	36341975	Gene Expr Patterns	WB,IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 23995-1-AP (FNDC5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 23995-1-AP (FNDC5 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).