

For Research Use Only

DR4 Polyclonal antibody, PBS Only

Catalog Number: 24063-1-PBS

Featured Product



Basic Information

Catalog Number:

24063-1-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG20927

GenBank Accession Number:

BC012866

GeneID (NCBI):

8797

UNIPROT ID:

O00220

Full Name:

tumor necrosis factor receptor superfamily, member 10a

Calculated MW:

50 kDa

Observed MW:

37-55 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human

Background Information

Tumor necrosis factor (TNF)-related apoptosis-inducing ligand (TRAIL) is a member of the TNF superfamily. TRAIL activates apoptosis through the death receptors DR4 (also known as TRAILR1 and TNFRSF10A) and DR5 (also known as TRAILR2, KILLER and TNFRSF10B). DR4 and DR5 are single-pass type I membrane proteins that contain intracellular death domains (DD) and upon activation mediate apoptotic signals (PMID: 11163110). Binding of TRAIL to DR4 or DR5 results in receptor oligomerization and recruitment of FAS-associated protein with death domain (FADD) and caspase 8 to form a functional death-inducing signalling complex (DISC). Upon DISC formation, caspase 8 is cleaved and activated, which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis (PMID: 18813321). DR4 or DR5 promotes the activation of NF-kappa-B and play an important role in inflammation (PMID: 9430227, 19434100).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24063-1-AP (DR4 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 24063-1-PBS in a different storage buffer formulation.



1×10^6 HeLa cells were intracellularly stained with 0.25 μ g DR4 Polyclonal antibody (24063-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00017). This data was developed using the same antibody clone with 24063-1-PBS in a different storage buffer formulation.