For Research Use Only

BBOF1 Polyclonal antibody

Catalog Number:24337-1-AP





Basic Information	Catalog Number: 24337-1-AP	GenBank Accession Number: BC 104981	Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 850 ug/ml by Nanodrop; Source: Rabbit	GenelD (NCBI): 80127	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
		UNIPROT ID: O8ND07	
		Full Name:	
	Isotype: IgG	chromosome 14 open reading frame 45	
	Immunogen Catalog Number: AG19302	Calculated MW: 529 aa, 62 kDa	
		Observed MW: 62 kDa	
Applications	Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse, rat	Positive Co	ntrols:
		WB : mouse testis tissue, TT cells, rat testis tissue IHC : mouse testis tissue,	
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen	
Background Information	Basal body orientation factor 1 (BBOF1), also known as CCDC 176, contains two coiled-coil domains and is evolutionarily conserved in vertebrates. Recently, Xenopus bbof1 has been characterized as a cilia-associated factor that is induced by the transcription factor foxj1 to mediate the alignment and orientation of cilia in epidermal multi-ciliated cells. BBOF1 is required for sperm flagellar axoneme maintenance and male fertility in mice. BBOF1 expression is elevated during meiosis, and it is accumulated in round spermatids. BBOF1 deletion has compromised effects on growth, survival, and spermatogenesis.		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	storage	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data





mouse testis tissue were subjected to SDS PAGE followed by western blot with 24337-1-AP (BBOF1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 24337-1-AP (BBOF1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 24337-1-AP (BBOF1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).