## For Research Use Only

## AMOT Polyclonal antibody

Catalog Number: 24550-1-AP

**Featured Product** 

2 Publications

BC130294



**Purification Method:** 

WB 1:1000-1:9000

IHC 1:50-1:500

IF 1:20-1:200

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:500-1:2000

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number:

GeneID (NCBI):

150ul, Concentration: 1000 µg/ml by 154796

Nanodrop; Full Name: Source: angiomotin Rabbit Calculated MW: 1084 aa, 118 kDa Isotype:

IgG Observed MW: Immunogen Catalog Number: 80 kDa, 130 kDa

AG19784

24550-1-AP

**Tested Applications:** 

IF, IHC, IP, WB, ELISA

**Cited Applications:** 

IF, WB

Species Specificity:

human **Cited Species:** 

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

IP: HEK-293 cells,

IHC: human colon cancer tissue,

WB: HEK-293 cells, HEK-293T

IF: HEK-293 cells,

Positive Controls:

**Background Information** 

Angiomotin belongs to the motin family of angiostatin binding protein. The encoded protein is expressed predominantly in endothelial cells of capillaries as well as larger vessels of the placenta where it may mediate the inhibitory effect of angiostatin on tube formation and the migration of endothelial cells during the formation of new blood vessels. Most abundant expression was found in placenta and skeletal muscle. AMOT has two isoforms with 

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Qiang Guo	32292505	Theranostics	WB
Diala Kantar	33707576	Sci Rep	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




WB result of AMOT antibody (24550-1-AP, 1:6000) with si-Control and si-AMOT transfected HEK293 cells




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 24550-1-AP (AMOT antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 24550-1-AP (AMOT antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 24550-1-AP (AMOT antibody) at dilution of 1:4500 incubated at room temperature for 1.5 hours.



IP result of anti-AMOT(IP:24550-1-AP, 4ug; Detection:24550-1-AP 1:1000) with HEK-293 cells lysate 1280 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using 24550-1-AP (AMOT antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).