For Research Use Only

SYNJ1 Polyclonal antibody

Catalog Number:24677-1-AP

4 Publications



Basic Information

Catalog Number: 24677-1-AP

GenBank Accession Number: BC136603

Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 700 µg/ml by Nanodrop and 293 µg/ml by Bradford Full Name:

8867

WB 1:1000-1:8000 IHC 1:50-1:500

IF 1:20-1:200

Purification Method:

method using BSA as the standard;

synaptojanin 1 Calculated MW:

1612 aa, 178 kDa

Isotype: IgG

Observed MW: 140 kDa

Immunogen Catalog Number:

AG20364

Rabbit

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

CoIP, IHC, IP, WB

Species Specificity: human, mouse, rat

Cited Species:

human, mouse

WB: mouse brain tissue, NIH/3T3 cells, rat brain tissue

IHC: human brain tissue, human skeletal muscle tissue, rat brain tissue, rat skeletal muscle tissue

IF: NIH/3T3 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Synaptojanin 1, a polyphosphoinositide phosphatase, is expressed as two major alternatively spliced isoforms of $145\,kDa~(SJ145)~and~170~kDa~(SJ170),~which~are~thought~to~have~pleiotropic~roles~in~endocytosis,~signaling~and~actin$ function. SJ145 is highly enriched in nerve terminals where it participates in clathrin-dependent synaptic vesicle recycling. SJ170, which differs from SJ145 by the presence of a carboxy-terminal extension, is the predominant isoform in developing neurons and is expressed in a variety of tissues. (PMID:10801423).

Notable Publications

Author	Pubmed ID	Journal	Application
Saifeldin N Shehata	30880224	Cell Signal	WB,IP
David Furkert	32783964	Cell Chem Biol	WB
Xinmiao Xie	38152048	Ren Fail	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 24677-1-AP (SYNJ1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 24677-1-AP (SYNJ1 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human brain tissue slide using 24677-1-AP (SYNJ1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 24677-1-AP (SYNJ1 Antibody) at dilution of 1:200 (under 40x lens).




Immunohistochemical analysis of paraffinembedded rat skeletal muscle tissue slide using 24677-1-AP (SYN)1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using 24677-1-AP (SYNJ1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded rat skeletal muscle tissue slide using 24677-1-AP (SYN)1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 24677-1-AP (SYNJ 1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 24677-1-AP (SYNJ1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).