

For Research Use Only

# MN1 Polyclonal antibody, PBS Only

Catalog Number: 24697-1-PBS

Featured Product



## Basic Information

**Catalog Number:**

24697-1-PBS

**Size:**

100ug, Concentration: 1 mg/ml by Nanodrop;

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG20344

**GenBank Accession Number:**

BC156879

**GeneID (NCBI):**

4330

**UNIPROT ID:**

Q10571

**Full Name:**

meningioma (disrupted in balanced translocation) 1

**Calculated MW:**

1320 aa, 136 kDa

**Observed MW:**

136 kDa

**Purification Method:**

Antigen affinity purification

## Applications

**Tested Applications:**

WB, IHC, Indirect ELISA

**Species Specificity:**

human

## Background Information

MN1 (Transcriptional activator MN1), which is mainly located in nucleus. Highest expression is observed in fetal brain and skeletal muscle, and adult skeletal muscle. MN1 protein can interact with Brg1/Brm related factor (BAF) complex containing Smarca4/Brg1 and stabilize it on chromatin, thus maintaining the expression of hematopoietic progenitor cell-like genes. Under normal physiological conditions, MN1 protein is mainly expressed in granulocyte monocyte progenitor cells (GMP) in hematopoietic system, which plays an important role in the development and function of hematopoietic cells, and it is involved in regulating cell proliferation, differentiation, apoptosis and embryonic development. MN1 protein is related to many diseases, especially in leukemia (PMID: 23049943). MN1 gene rearrangements such as t(12; 22)(p13; Q11) can produce MN1-TEL fusion protein, which combines the transcriptional activation domain of MN1 and the DNA binding domain of TEL(ETV6), and can stably occupy the TEL recognition sequence, hindering the combination of normal transcription regulatory factors, thus leading to leukemia. Overexpression of MN1 gene has also been proved to be one of the signs of poor prognosis in patients with acute myeloid leukemia (AML), and its expression level is high in AML patients with normal karyotype. The molecular weight of MN1 is 136 kDa.

## Storage

**Storage:**

Store at -80°C.

**Storage Buffer:**

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24697-1-AP (MN1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 24697-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 24697-1-AP (MN1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 24697-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 24697-1-AP (MN1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 24697-1-PBS in a different storage buffer formulation.