For Research Use Only

FAM172A Polyclonal antibody

Catalog Number:24732-1-AP



Purification Method:

WB 1:1000-1:6000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

 24732-1-AP
 BC020584

 Size:
 GeneID (NCBI):

 150ul , Concentration: 700 ug/ml by
 83989

Nanodrop; UNIPROT ID:
Source: Q8WUF8
Rabbit Full Name:

Isotype: family with sequence similarity 172, IgG member A

IgGmember AImmunogen Catalog Number:Calculated MW:AG20437416 aa, 48 kDa

Observed MW: 50-55 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse colon tissue, rat colon tissue

IHC: human liver tissue,

Background Information

Family with sequence similarity 172 member A (FAM172A), cloned from human aortic tissues, is subsequently observed in human endothelial and vascular smooth muscle cells, and macrophages. As the imbalance of the cell cycle and apoptosis is closely related to the key characteristics of malignant tumors, FAM172A dysregulation is a potential contributing factor in cell transformation. Recently, FAM172A has been reported to take part in the development of several cancers, such as human liver, colorectal and papillary thyroid cancers. Moreover, FAM172A was upregulated by high glucose in a concentration- and time-dependent manner which indicated that FAM172A may be involved in the pathogenesis of diabetic related diseases. FAM172A has 4 isoforms with the molecular mass of 34-36, 43 and 48 kDa. This antibody was detected at 50-55 kDa on SDS-PAGE. (PMID: 31988090, 26637224)

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24732-1-AP (FAM172A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 24732-1-AP (FAM172A Antibody) at dilution of 1:200 (under 40x lens).