For Research Use Only

BAALC Polyclonal antibody

Catalog Number:24997-1-AP 1 Publications

Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 24997-1-AP	GenBank Accession Number: BC011517	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	Nanodrop and 267 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG21850	79870 UNIPROT ID:	WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total
		Q8WXS3 Full Name: brain and acute leukemia, cytoplasmic Calculated MW:	protein lysate
			IHC 1:50-1:500 IF/ICC 1:50-1:500
		19 kDa	
		Observed MW: 19-25 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, IF/ICC, IP, ELISA	WB: H	IEK-293 cells, A375 cells, mouse brain tissue
	Cited Applications: IHC	IP : m	ouse brain tissue,
	Species Specificity:	IHC : I	mouse brain tissue,
	human, mouse	IF/ICC : HEK-293 cells,	
	Cited Species: human		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen	
Background Information	cells. Expression is found both in norr	cytoplasm and membrane. BA mal early progenitor cells and	.C, is a 180 amino acid protein, which interact ALC is expressed in neural and hematopoiet in the most immature type of blasts in acute I synaptic role at the postsynaptic lipid rafts b
Background Information	cells. Expression is found both in norr leukemia but not in mature hematop interacting with CAMK2A.	cytoplasm and membrane. BA mal early progenitor cells and	ALC is expressed in neural and hematopoiet in the most immature type of blasts in acute


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 24997-1-AP (BAALC Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-BAALC (IP:24997-1-AP, 4ug; Detection:24997-1-AP 1:300) with mouse brain tissue lysate 3000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using 24997-1-AP (BAALC antibody) at dilution of 1:50 and Alexa Fluor 488conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 24997-1-AP (BAALC antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).